



Ministry of Health & Family Welfare
Government of India

INTEGRATED DISEASE SURVEILLANCE PROJECT (IDSP)

NON-COMMUNICABLE DISEASE RISK FACTORS SURVEY

2007-08

Andhra Pradesh



Indian Institute of Health & Family Welfare
Hyderabad
(State Survey Agency)

National Institute of Epidemiology
Chennai
(Regional Resource Centre)

National Institute of Medical Statistics
New Delhi
(National Nodal Agency)

National Institute of Communicable Diseases
New Delhi
(IDSP Central Surveillance Unit)

Indian Council of Medical Research
New Delhi
(National Implementing Agency)

Funded by the World Bank

2009

Suggested Citation

National Institute of Medical Statistics, Indian Council of Medical Research (ICMR), 2009, IDSP Non-Communicable Disease Risk Factors Survey, Andhra Pradesh, 2007-08. National Institute of Medical Statistics and Division of Non-Communicable Diseases, Indian Council of Medical Research, New Delhi, India.

Contents

Page No

Foreword	v
Preface	vii
Acknowledgements	ix
Definitions	xi
Acronyms	xiii
Tables	xv
Figures	xvii
Fact Sheet	xix
Executive Summary	xxi
1. Introduction	1
1.1 Background of the Survey	1
1.2 Objectives	1
1.3 Non communicable Disease(NCD) Risk Factors	1
1.4 Health Profile of the State	2
1.5 Survey Design and Implementation	3
1.6 Survey Instruments	5
1.7 Training	6
1.8 Data Processing and Analysis	7
1.9 Quality Control Measures	7
2. Background Characteristics of the Respondents and Households	9
2.1 Household Characteristics	9
2.2 Age and Sex Composition	10
2.3 Education Level	10
2.4 Marital Status	10
2.5 Occupation	11
3. Behavioural Risk Factors	13
3.1 Tobacco Smoking and Chewing	13
3.2 Alcohol Consumption	17
3.3 Fruits and Vegetables Consumption	19
3.4 Physical Activity	21
3.5 Socio-Demographic Pattern	23

4.	Hypertension and Diabetes	29
4.1	Hypertension	29
4.2	Socio-Demographic Pattern of Hypertension	31
4.3	Diabetes	32
5.	Physical Measurements	33
5.1	Weight	33
5.2	Height	33
5.3	Body Mass Index (BMI)	33
5.4	Waist Circumference	33
5.5	Socio-Demographic Pattern of Overweight	35
6.	Summary and Conclusions	37
	Reference	39
	Appendix A Sample Weights	41
	Appendix B Household/Individual Questionnaire	44
	Appendix C Technical Support and Human Resource	58



डॉ विश्व मोहन कटोच

एम डी, एक एन ए एससी, एक ए एम एस, एक ए एसी, एक एन ए

सचिव, भारत सरकार

(स्वास्थ्य अनुसंधान विभाग)

स्वास्थ्य एवं परिवार कल्याण मंत्रालय एवं

महानिदेशक, आई सी एम आर

Dr. Vishwa Mohan Katoch

MD, FNASc, FAMS, FASc, FNA

Secretary to the Government of India

(Department of Health Research)

Ministry of Health & Family Welfare &

Director-General, ICMR



भारतीय आयुर्विज्ञान अनुसंधान परिषद

(स्वास्थ्य अनुसंधान विभाग)

स्वास्थ्य एवं परिवार कल्याण मंत्रालय

वी. रामलिंगस्वामी भवन, अंसारी नगर

नई दिल्ली - 110 029 (भारत)

Indian Council of Medical Research

(Department of Health Research)

Ministry of Health & Family Welfare

V. Ramalingaswami Bhavan, Ansari Nagar

New Delhi - 110 029 (INDIA)

Foreword

Globally, non-communicable diseases (NCDs) are the major cause of morbidity and mortality. According to WHO Report 2004, they account for almost 60% of deaths and 47% of the global burden of disease. In India, estimated deaths due to non-communicable diseases were double than those from communicable diseases. A progressive rise in the disease pattern of NCD foretells a serious public health issue. The major risk factors for non-communicable diseases are tobacco and alcohol abuse, a sedentary lifestyle, and an unhealthy diet. It is believed that about half of non-communicable disease-related premature deaths could be prevented through healthy diet, regular physical activity and by avoiding tobacco and alcohol.

Envisaging the magnitude of the public health problem of chronic diseases, the Government of India through National Institute of Communicable Disease, MoHFW and Indian Council of Medical Research initiated NCD risk factors survey, phase-I in seven states of India. It is a well planned large community based survey providing state wise estimates of major NCD risk factors in different strata of population. It is needless to mention that the estimated NCD risk factors are important input for targeted prevention of NCD and effective health care planning. The National Technical Advisory Committee (NTAC) and National Monitoring Committee constituted by MoHFW, provided valuable technical guidance and support to complete the study.

The Indian Council of Medical Research through its Division of Non-communicable Diseases has implemented the study with all the partners including the National Institute of Medical Statistics as a National Nodal Agency, Regional Resource Centers and State Survey Agencies.

I congratulate the Team for successfully completing the survey and bringing out Phase-I report of NCD Risk Factors which would be of immense use for prevention and control of non-communicable diseases.

(Dr. V.M. Katoch)

डॉ बेला शाह एम.डी.
वैज्ञानिक 'जी' एवं प्रमुख
असंचारी रोग प्रभाग

Dr. Bela Shah MD
Scientist 'G' & Head
Division of Non-Communicable Diseases



भारतीय आयुर्विज्ञान अनुसंधान परिषद
स्वास्थ्य अनुसंधान विभाग
स्वास्थ्य एवं परिवार कल्याण मंत्रालय
वी. रामलिंगस्वामी, अंसारी नगर
नई दिल्ली-110 029 (भारत)

Indian Council of Medical Research
Department of Health Research
Ministry of Health & Family Welfare
V. Ramalingaswamy Bhawan
Ansari Nagar, New Delhi - 110 029 (INDIA)

Preface

Non-Communicable Diseases (NCD) account for a large proportion of morbidity and mortality amongst the adult population of our country. The high prevalence of major risk factors viz. tobacco and alcohol consumption, inappropriate diet, physical inactivity, high blood pressure, high blood glucose and dyslipidemias are driving the epidemic of NCDs. The Division of Non-Communicable Diseases at ICMR was identified as the nodal point for surveillance of NCDs and their risk factors by the World Health Organization, and multi-site studies helped us in developing a sound strategy for NCD risk factor surveys at the national level under IDSP. The survey methodology developed by ICMR was incorporated by IDSP into the overall survey protocol provided to ICMR for implementation.

The Indian Council of Medical Research signed a Memorandum of Understanding (MOU) with IDSP for the standardization and quality assurance of the NCD risk factor surveys under the World Bank funded IDSP on behalf of Ministry of Health, Govt. of India. As per IDSP plan, these surveys were to be carried out in three phases so as to cover all States and UTs of the country. In the present Phase I, the State based estimates of the risk factors in seven States (Andhra Pradesh, Kerala, Madhya Pradesh, Maharashtra, Mizoram, Tamil Nadu and Uttarakhand) were arrived at through the IDSP identified seven State Survey Agencies, five Regional Research Centers and a National Nodal Agency under the overall guidance and supervision of ICMR Headquarters through the National Technical Advisory Committee. I am grateful to the Director General, ICMR for supporting the Division of Noncommunicable Diseases ICMR to implement the surveys. The untiring effort of our partner agencies is commendable and is duly acknowledged.

This report marks an important milestone in surveillance activities for NCDs in the country. The results would be useful for planning and monitoring an effective response in a coordinated manner by the Government. It should also stimulate further analysis and research in the area.

(Dr. Bela Shah)

Acknowledgements

The National Institute of Medical Statistics was identified as National Nodal Agency (NNA) to conduct the IDSP-NCD Risk Factors Survey Phase-I in India. The survey was carried out with the joint efforts of all partner organizations including Division of Non-Communicable Diseases of Indian Council of Medical Research as the implementing agency; and Center for Community Medicine, All India Institute of Medical Sciences, New Delhi; Regional Medical Research Centre for Tribal, Jabalpur; National Institute of Epidemiology, Chennai; Sri Chitra Tirunal Institute of Medical Sciences and Technology, Thiruvananthapuram; Regional Medical Research Centre, Dibrugarh as Regional Resource Centers; Department of Community Medicine of Chhatrapati Shahuji Maharaj Medical University, Lucknow; Government Medical College, Nagpur; Pune Health Care Management and Research Centre, Pune; Indian Institute of Health and Family Welfare, Hyderabad; Madras Diabetes Research Foundation, Chennai; Clinical Epidemiology Unit, Medical College, Thiruvananthapuram; Regional Institute of Medical Sciences, Imphal, Manipur as State Survey Agencies respectively.

We sincerely acknowledge the Ministry of Health and Family Welfare (MoHFW), Government of India for granting us responsibility of conducting the IDSP NCD Risk Factors Survey Phase-I in India. We acknowledge the World Bank for providing financial support to conduct the Phase-I survey in seven states. We gratefully acknowledge the technical support and valuable guidance provided by Dr. N.K. Ganguly, Chairman and all members of National Technical Advisory Committee (NTAC) and Dr. Shiv Lal, Special Director General Health Services, Director NICD and all the members of National Monitoring Committee. Thanks go to Dr. G. Ramana and J. Gowrinath Sastry from World Bank; Dr. Cherian Varghese, WHO; Dr. D. Bachani, Dr. R.L. Ichhpujani, Dr. A.C. Dhariwal, Dr. Shah Hossain and Dr. Pradeep Khasnobis from IDSP Central Surveillance Unit, NICD for their support in undertaking the survey. We are grateful to Dr. L.M. Nath and Dr. K. Anand, AIIMS, New Delhi and Dr. B.N. Bhattacharya, Indian Statistical Institute, Kolkatta for their technical guidance and review of the reports. We are extremely thankful to Dr. Bela Shah and her colleagues Dr. D.K. Shukla and Dr. Prashant Mathur at ICMR for providing leadership to implement the survey.

The team of NIMS including Dr. H.K. Chaturvedi, Dr. D. Sahu, Dr. Tulsi Adhikari, Dr. Atul Juneja, Mr. Jiten Kumar Singh and all other supporting staff involved in the study deserve appreciation and acknowledgement. We are grateful to National Institute of Epidemiology, Chennai being the Regional Resource Center for Andhra Pradesh and Indian Institute of Health and Family Welfare, Hyderabad involved as State Survey Agency for supervising, data collection and data entry of survey in Andhra Pradesh.

The hard work of all the field investigators, field supervisors and data entry operators are highly appreciable and acknowledged. Last but not the least, I express my heartiest thanks to all the respondents and other peoples including local health administrators of districts and state who helped in completing the survey.

National Institute of Medical Statistics
ICMR, New Delhi



(Arvind Pandey)
Director

Definitions

Current Smoker / Smokeless Tobacco User: Some one who at the time of the survey, smokes / uses tobacco in any form either daily or occasionally.

Current Daily Smoker / Smokeless Tobacco User: Some one who smokes / uses tobacco everyday with rare exceptions such as not on days of religious fasting or during acute illness.

Past- Daily Smokers / Smokeless Tobacco User: These are those individuals who were smoking daily in past, but have not smoked ever in one year preceding the survey.

Non-Smoker / Never Used Smokeless Tobacco: These are those individuals who have never smoked / used smokeless tobacco in the lifetime.

Current Drinker: Those who consumed one or more than one drink of any alcohol in the year preceding the survey.

Former Drinker: Those who have consumed alcohol but those who did not consume one or more drink during the year preceding the survey.

Lifetime Abstainer: Those who have never consumed one or more drink of any type of alcohol in lifetime.

High Risk Drinker (Binge Drinker): Those who drink more than 5 (for women 4) standard drinks on any single day.

Standard Drink: It is defined as any standard drink with net alcohol content of 10 gm ethanol.

Standard Serving: One standard serving of fruits and vegetables is equivalent to 80 grams, translated into different units of cups depending on type of vegetables and fruits.

Metabolic Equivalent (MET): MET is the ratio of a person's working metabolic rate relative to the resting metabolic rate. One MET is defined as the energy cost

of sitting quietly, and is equivalent to a caloric consumption of 1 kcal/kg/hour. It is estimated that, compared to sitting quietly, a person's caloric consumption is four times as high when being moderately active, and eight times as high when being vigorously active.

Central Obesity: Central obesity (measured as waist circumference or waist to hip ratio) is more strongly associated with coronary heart disease than BMI. Waist measurement is taken at the level of mid point between the inferior margin of the rib and crest of the ileum in the mid-axillary plane, using a non-stretchable tape, without clothing, that is, directly over the skin (or over light clothing). A cut-off level of 102 centimeters in males and 88 centimeters in females have been recommended for developed countries (ATP 3 Guidelines), however, much lower cut-off levels are appropriate for Indians of 90 centimeters in males and 80 centimeters in females (South Asia Pacific Guidelines).

Hypertensive Stage I: The upper and lower limit of the systolic and diastolic blood pressure for hypertensive stage I is 140-159 mm Hg systolic or 90-99 mm Hg for diastolic.

Hypertensive Stage II: The upper and lower limit of the systolic and diastolic blood pressure for hypertensive stage II is ≥ 160 mm Hg systolic or ≥ 100 mm Hg for diastolic.

Under Weight: The person with BMI less than 18.5 kg/m² is categorized as under weight.

Normal Weight: The person whose BMI is between 18.5 to 24.9 kg/m² is categorized as normal weight.

Over Weight: The person whose BMI is 25 kg/m² or more is categorized as over weight.

Acronyms

AYUSH	Ayurveda, Unani, Shidha and Homeopathy
BMI	Body Mass Index
BP	Blood Pressure
CEB	Census Enumeration Block
DHO	District Health Officer
ICMR	Indian Council of Medical Research
IDSP	Integrated Disease Surveillance Project
IIHFW	Indian Institute of Health and Family Welfare
LPG	Liquid Petroleum Gas
MET	Metabolic Equivalent
NCD	Non-communicable Diseases
NICD	National Institute of Communicable Diseases
NIE	National Institute of Epidemiology
NIMS	National Institute of Medical Statistics
NMC	National Monitoring Committee
NNA	National Nodal Agency
NTAC	National Technical Advisory Committee
PSU	Primary Sampling Unit
RRC	Regional Resource Centre
SSA	State Survey Agency
WC	Waist Circumference
WHO	World Health Organization

Tables

	Page No.	
Table 1.1	Demographic and Socioeconomic profile of Andhra Pradesh as compared to India	3
Table 1.2	Health Infrastructure, Human Resource available and Health Expenditure	3
Table 1.3	Sample coverage and response rate of household, step-1 and step-2 individual response rate by place of residence, Andhra Pradesh, 2007- 08	5
Table 2.1	Percentage distribution of households in rural and urban area according to the background Characteristics, Andhra Pradesh, 2007-08	9
Table 2.2	Age and sex wise distribution of respondents by residence (un-weighted) Andhra Pradesh, 2007-08	10
Table 2.3	Percentage of respondents according to background characteristics, sex and place of residence, Andhra Pradesh, 2007-08.	11
Table 3.1.1	Percentage of respondents classified by smoking status across sex and place of residence, Andhra Pradesh, 2007- 08	13
Table 3.1.2	Mean number of tobacco products smoked (per day) by daily smokers according to sex and place of residence, Andhra Pradesh, 2007-08.	14
Table 3.1.3	Mean age of initiation, age at stopped smoking and percentage of respondents (non-smokers) exposed to tobacco smoke by sex and place of residence, Andhra Pradesh, 2007- 08	14
Table 3.1.4	Percentage of smokeless tobacco users by sex and place of residence, Andhra Pradesh, 2007-08	15
Table 3.1.5	Mean frequency of smokeless tobacco use (per day) by daily smokeless tobacco users according to sex and place of residence, Andhra Pradesh, 2007-08	15
Table 3.1.6	Mean age of initiation, age at stopped smokeless tobacco use by daily smokeless tobacco users according to sex and place of residence, Andhra Pradesh, 2007-08	16
Table 3.1.7	Percentage of tobacco users by sex and place of residence, Andhra Pradesh, 2007-08	16
Table 3.2.1	Percentage of Alcohol consumption by sex and place of residence, Andhra Pradesh, 2007- 08	17
Table 3.2.2	Percentage of drinkers (past 12 months) according to frequency of drinking, mean number of standard drinks per day and pattern of drinking in the last seven days by sex and place of residence, Andhra Pradesh, 2007-08	17
Table 3.2.3	Mean age of initiation of alcohol use by sex and place of residence, Andhra Pradesh, 2007-08	18
Table 3.2.4	Percentage of current daily smokers, daily smokeless tobacco users and current drinkers across age, education and occupation, Andhra Pradesh, 2007-08	19
Table 3.3.1	Mean number of days in a week fruits and vegetables consumed by the respondents according to sex and place of residence, Andhra Pradesh,2007-08	19
Table 3.3.2	Mean number of servings of fruits, vegetables consumed in one particular day by sex and place of residence, Andhra Pradesh, 2007-08	20

Table 3.3.3	Percentage of respondents according to the intake of specific food items by sex and place of residence, Andhra Pradesh, 2007-08.	20
Table 3.3.4	Percentage of households according to type of oil consumption, Andhra Pradesh, 2007-08	21
Table 3.4.1	Mean time spent (in METminutes) on physical activity per day by sex and residence, Andhra Pradesh, 2007-08	21
Table 3.4.2	Percentage of respondents classified in three categories of total physical activity per day (P & 95% CI) by sex and place of residence, Andhra Pradesh, 2007-08	22
Table 3.4.3	Percentage of respondents (with 95% Confidence Interval) according to three categories of total physical activity by age group and sex, Andhra Pradesh, 2007-08	22
Table 3.4.4	Sex wise percentage of respondents classified according to total time spent in sedentary activity per day by residence, Andhra Pradesh, 2007-08	23
Table 3.5.1	Percentage of respondents in the category of some high risk factors of NCD (current daily smokers, daily smokeless tobacco user, current drinkers, low fruits and vegetables intake and low physical activity) across age, education, occupation and sex, urban, Andhra Pradesh, 2007- 08	24
Table 3.5.2	Percentage of respondents in the category of some high risk factors of NCD (current daily smokers, daily smokeless tobacco user, current drinkers, low fruits and vegetables intake and low physical activity) across age, education, occupation and sex, rural, Andhra Pradesh, 2007- 08	26
Table 3.5.3	Percentage of respondents in the category of some high risk factors of NCD (current daily smokers, daily smokeless tobacco user, current drinkers, low fruits and vegetables intake and low physical activity) across age, education, occupation and sex, combined, Andhra Pradesh, 2007- 08	27
Table 4.1.1	Percentage of respondents with history of raised blood pressure, treatment and life style modification advised, seeking consultation and treatment from an AYSUH practitioner by sex and place of residence, Andhra Pradesh, 2007-08	29
Table 4.1.2	Mean Systolic and Diastolic blood pressure by sex and place of residence, Andhra Pradesh, 2007-08	30
Table 4.1.3	Categories of Hypertension	30
Table 4.1.4	Percentage of respondents according to category of hypertension by sex and place of residence, Andhra Pradesh, 2007-08	30
Table 4.2	Percentage of respondents in the category of stage I & stage II hypertension across age, education, occupation and by sex and residence, Andhra Pradesh, 2007- 08	31
Table 4.3	Percentage of respondents with history of raised blood sugar, treatment and life style modification advised, seeking consultation and treatment from an AYSUH practitioner by sex and place of residence, Andhra Pradesh, 2007-08	32
Table 5.1	Categories of BMI	33
Table 5.2	Mean value for body mass index (BMI), height, weight and waist circumference by sex and place of residence, Andhra Pradesh, 2007-08	34
Table 5.3	Percentage of respondents according to BMI categories by sex and place of residence, Andhra Pradesh, 2007-08	34
Table 5.4	Percentage of respondents in the category of overweight (Grade I, II & III) across age, education, occupation and by sex and residence, Andhra Pradesh, 2007- 08	35

Figures

	Page No.	
Figure 1	Current smokers (%) by sex and residence	xii
Figure 2	Current daily smokeless tobacco users (%) by sex and residence	xii
Figure 3	Alcohol consumption (%) by sex and residence	xxiii
Figure 4	Less than five servings of fruits and vegetables consumption (%) by sex and residence	xxiii
Figure 5	Major oil consumption among households (%) by residence	xxiii
Figure 6	Mean time spent on physical activity per day (minutes) by sex and residence	xxiii
Figure 7	Low physical activity of respondents (%) by sex and residence	xxiv
Figure 8	Stage I & II hypertension (%) by sex and residence	xxiv
Figure 9	History of raised blood sugar (%) by sex and residence	xxiv
Figure 10	Overweight respondents (%) by sex and residence in Andhra Pradesh	xxiv

IDSP-NCD Risk Factor Survey

Fact Sheet - Andhra Pradesh

Population		<i>Any form of tobacco use</i>	24
Household covered	4905	Male	39
Individual covered	6218	Female	8
Household Characteristics(%)		Mean age of Initiation (in years)	
Religion		<i>Smoking</i>	19
Hindu	82	Male	19
Muslim	8	Female	14
Access to piped drinking water	75	<i>Smokeless tobacco</i>	20
Urban	94	Male	20
Rural	68	Female	23
Sanitation		Alcohol Consumption	
Flush Toilet	47	<i>Consumed alcohol (last 30 days)</i>	14
Urban	89	Male	27
Rural	32	Female	2
Source of Lighting		<i>Consumed alcohol (last 12 Months)</i>	20
Electricity	93	Male	37
Urban	99	Female	3
Rural	91	Consumed alcohol (in last 7 days)	
Type of House		<i>Binge Drinkers</i>	23
Pucca house	54	Male	24
Urban	70	Female	11
Rural	49	Mean age of Initiation Alcohol (in years)	
Kachha house	19	Male	20
Urban	5	Female	15
Rural	24	Fruits and Vegetables Consumed	
Fuel Use for Cooking		<i>Less than five servings per day</i>	88
LPG / Gas	31	Urban	86
Urban	72	Rural	90
Rural	17	Physical Activity	
Wood	65	<i>Low physical activity</i>	68
Urban	21	Urban	78
Rural	81	Rural	64
Separate Kitchen		Hypertension	
Urban	60	<i>Pre hypertension</i>	43
Rural	38	Urban	46
Own Agricultural land		Rural	42
Urban	7	<i>Stage I and II hypertension</i>	17
Rural	37	Urban	16
Individual Characteristics(%)		Rural	17
Education -Illiterate	45	Physical Measurement - BMI	
Urban (Male)	17	<i>Under weight</i>	25
Urban (Female)	36	Urban	17
Rural (Male)	41	Rural	28
Rural (Female)	64	<i>Over weight (grade I,II,III)</i>	19
Behavioural Information(%)		Urban	31
Current Tobacco Users		Rural	15
Smokers	18	<i>Central obesity</i>	21
Male	32	Urban	30
Female	4	Rural	17
Smokeless tobacco users	9		
Male	14		
Female	5		

Executive Summary

Introduction

The Government of India through the Ministry of Health & Family Welfare (MOHFW) initiated a decentralized, state based Integrated Disease Surveillance Project (IDSP) in the country with the assistance of the World Bank in the year 2004. The component of non communicable disease surveillance planned periodic community based surveys of population aged 15-64 to provide data on the risk factors. It is in line to help the state health administrators to plan strategies for the control of non communicable diseases by modifying the risk factors. All Indian states were proposed to be surveyed in a phased manner under the project. The first phase of the survey included seven states namely Andhra Pradesh, Kerala, Madhya Pradesh, Maharashtra, Mizoram, Tamil Nadu and Uttarakhand.

The overall objective of the NCD risk factors survey was to improve the information available to the Government health services and care providers on a set of high-priority risk factors, with a view to improve the quality health care and services. The survey also aimed to establish the baseline database of NCD risk factors needed to monitor trends in population health behavior and risk factors for chronic diseases over time. This would provide evidence for evolving strategies and interventions for identified risk factors in the community to reduce the burden of non-communicable diseases.

A National Technical Advisory Committee was constituted to provide the technical guidance to the survey including taking care of certain administrative and logistic difficulties and the National Monitoring Committee for monitoring the overall progress of the project. Indian Council of Medical Research through the division of non communicable diseases, was the implementing agency while the National Institute of Medical Statistics (NIMS) was appointed as the National Nodal Agency (NNA) for coordinating the survey; the National Institute of Epidemiology, Chennai as a Regional Resource Centre (RRC) for monitoring the quality of data collection and technical support to State Survey Agency (SSA) for the state of Andhra Pradesh.

Survey Methodology

WHO STEPS methodology for NCD Risk Factor Surveillance has been adopted for the survey after carrying out suitable modifications, based on a multi-site ICMR-WHO collaborative initiative for NCD risk factor surveillance¹. The survey was designed to provide prevalence estimates of risk factors for each 10 years age group (15-24 through 55-64) by sex and place of residence (urban/rural). The survey used uniform sample design, bilingual schedules (English and Telegu in case of Andhra Pradesh), field protocol for data collection and physical measurements to facilitate comparability across states and also to ensure high quality data. For the present survey, appropriate sampling weights for households were used for urban and rural areas of the state. From each selected household one member aged 15-54 was selected using the KISH Method and all members aged 55-64 were selected. Such post stratification was used for improvement of efficiency of the estimators. Post stratification weights for individuals were constructed using the state age distributions for both sexes, which are available on the population level.

Two types of questionnaire - one at household level and another for individual level were used for the survey. At household level, information was elicited on religion, household facilities, ownership of agricultural land and livestock, and possession of durable goods for each selected household. The Individual questionnaire collected the information from all the selected eligible household individuals regarding demographic, behavioral and physical measurements. The individual questionnaire was divided into two segments based on WHO STEP methodology. The first section (Step 1) collected the demographic information of individuals including age, sex, marital status, education, and occupation. In the behavioural information section, information about tobacco use, alcohol consumption, diet, physical activity, history of raised blood pressure and history of diabetes were collected. In the second section (Step 2), physical measurements of individual such as height, weight, waist circumference (not measured for pregnant women), blood pressure, and pulse rate were recorded.

Characteristics of Survey Population

A total of 5000 households were contacted in urban and rural area of Andhra Pradesh. Among them 95 households refused to participate in the survey. The overall response for the survey was therefore over 98 percent. More than four-fifth (82%) of the households was Hindu, about 11% was Christian and about 8% was Muslim. Ninety percent of the households used drinking water from a piped or hand pump. All the households had flush or pit toilet facility. Ninety-three percent of households had electricity. LPG was a major source of cooking fuel in urban area and wood was main source of fuel in rural area. As envisaged, 37% of the households in rural possessed agricultural land, where as it was only 7% in case of urban.

Almost half of the respondents (45%) were illiterate. It was pronounced in rural area where almost 2/3rd (64%) of females and 41% of males were illiterate. The results emphasize the need of taking female literacy program of the Government. About three-quarter (72%) of the respondents were currently married. About 50% of females were engaged in domestic work. Majority of the males in rural area were engaged in agricultural and manual work where as in urban areas they were mostly engaged in manual work and executive/business positions.

BEHAVIOURAL RISK FACTORS FOR NCD

Tobacco Smoking

As per the WHO STEPS guidelines, the smokers are categorized into three categories *Current Smokers*, *Current Daily Smokers*, *Past Daily Smokers* and those who have never smoked in lifetime are classified as *Non-Smokers*. About one fifth of respondents (32 percent men and about 4 percent women) in Andhra Pradesh were current smokers.




Figure 1. Current smokers (%) by sex and residence

The urban and rural prevalence were 13 % and 20% respectively for the current smokers. The mean number of smoking *beedis* and manufactured cigarette in a day was 5 and 3 respectively. There was marked difference between men and women in the mean frequency of any type of smoking. The average age of onset of smoking was around 21 years among young rural respondents as compared to 20 years among young urban respondents aged 15-34 years. The mean age of cessation of smoking for all those who stopped smoking was 27 years. Among non-smokers one fifth of respondents were exposed to tobacco smoke at home or work.




Figure 2. Current daily smokeless tobacco users (%) by sex and residence

Nine percent of the respondents (14% of males and 5% of females) were current users of smokeless tobacco. The mean frequency of chewing tobacco in a day was less than one for tobacco chewer. The mean frequency of chewing pan with tobacco was one for both, male and female. Snuff by mouth/nose was around 2. The mean age of initiation of smokeless tobacco use was 20 years for the young respondents aged 15-34 years and it was 25 years for the respondents in the age group of 35-64 years. The over all mean age of quitting smokeless tobacco was 27 years. About 24% of the respondents were using tobacco in either form (smoking or smokeless) whereas about 2% were using tobacco in both the forms.

Alcohol Consumption

In the survey, 20% of the respondents report to have consumed alcohol in past 12 months and 14% consumed in last 30 days preceding the survey. Only 3% of respondents were past drinkers. The habit was higher among men with 37% consuming alcohol in past 12 months as compared to only 3% among women. The average number of drinks on a drinking day was 3 drinks. About 15% of the respondents had 20 or more drinks in

the last 7 days. The mean age of initiation of alcohol consumption regularly was 20 years for the respondents in the age group 15-34 years and 25 years for the respondents in the age group of 35-64 years. Twenty seven percent of men and only 2 percent of women report to have consumed alcohol in past 30 days. The percentage of current drinkers was high for respondents whose main occupation is agriculture or manual work.




Figure 3. Alcohol consumption (%) by sex and residence

Fruits and Vegetables Consumption

In a week, people consumed vegetables 5 days and fruits only 2.5 days. The mean number of days when fruits were consumed was higher in urban population (2.5 days) as compared to that of rural (1.6 days). Only one tenth (12%) of population consumed five or more servings of fruits and vegetables per day.




Figure 4. Less than 5 servings of fruits & vegetables (%) by sex and residence

Food and Oil Consumption

In respect of consumption of specific food habits, about 70% population consumed eggs, about 50% consumed chicken, about 28% consumed fish and 34%

consumed red meat at least once a week. Cheese/butter was consumed daily by 3% of the population.




Figure 5. Major oil consumption among households (%) by residence

Most common edible oil used for cooking in Andhra Pradesh was Palm oil (49%), followed by groundnut oil (29%), and sunflower oil (19%).

Physical Activity

The lack of physical activity leads to obesity, hyperlipidemia, diabetes mellitus, hypertension, and coronary heart disease. In this respect survey finds that in Andhra Pradesh, the mean time spent in different sub groups on work related physical activity ranged between 188 to 246 minutes per day. The mean duration of physical activity was 269 minutes per day. Most of the time spent was related to work only. Around 31 minutes per day was spent for travel activities.




Figure 6. Mean time spent on physical activity per day (minutes) by sex and residence

As per the WHO guidelines, the total physical activity of the individual has been categorized as low, medium and high. About 68% of respondents report low level of physical activity, while 28% and 4% of respondents report medium and high level of activity,




Figure 7. Low physical activity of respondents (%) by sex and residence

respectively. Majority of respondents spent 2 to 3 hours in sedentary activities (33%).

HYPERTENSION AND DIABETES

Hypertension

The blood pressure is an important determinant of risk of cerebrovascular and ischemic heart diseases, congestive cardiac failure and renal failure. In the survey, 8% respondents report to have been diagnosed as hypertensive by health professionals (7% for males and 8% for females; 10% in urban and 6% in rural population). Among those who were diagnosed with hypertension, 67% were on prescribed drugs, 71% were advised dietary modification, and 8% consulted AYUSH practitioner of which three-fourth were taking treatment from the same. The survey also carried out measurement of blood pressure as a part of step two of individual questionnaire. The mean systolic blood pressure in the population was around 123 mm Hg and mean diastolic blood pressure was 77 mm Hg. By categories of hypertension, 36% report to be normal, 43% in pre-hypertension, 15% in Stage I hypertension and 6% in Stage II hypertension.




Figure 8. Stage I & II hypertension (%) by sex & residence

Diabetes

Diabetes mellitus is an important marker of risk for the arterial disease of the coronary, cerebral and peripheral arterial trees, and for micro vascular disease leading to blindness and renal failure. The survey also included information on history of diabetes. Around 2% of the respondents report to have history of raised blood sugar of which 16% were taking insulin and 86% were on oral drugs. A large number of them were advised life style modification such as diet modification, reducing weight and increasing physical activity. About 8% (of the diagnosed) had consulted AYUSH practitioners for the elevated blood sugar levels and majority of them were taking the treatment from the system.




Figure 9. History of raised blood sugar (%) by sex and residence

PHYSICAL MEASUREMENTS

Body Mass Index (BMI)

Worldwide researches have shown that there is a strong association between BMI and health risk. On the




Figure 10. Overweight respondents (%) by sex and residence in Andhra Pradesh

other hand low BMI is an indicator of risk to health, often being associated with tobacco, alcohol use and drug addiction. The survey recorded on height, weight and waist circumference. The mean BMI was around 22 kg/m² with mean height 157.6 cm (163.5 cm for men, 151.6 cm for women) and mean weight 54.0 kg (57.9 kg for men and 49.7 kg for women). According to the survey, 25% of the respondents were under weight and about 19% were overweight which was 30% in urban and 17% in rural population. Overall 56% of the respondents were in the normal category of BMI.

SOCIO-DEMOGRAPHIC DIFFERENTIALS

Tobacco is mainly used either in the form of smoking or non-smoking. The prevalence of smoking was high among urban as well as rural male population. The increasing pattern of prevalence of smoking was recorded with increasing age group of respondents. But, it was declining with increasing level of education. Prevalence of smoking among female respondents was very low compare with males across all the socio-demographic categories, which shows gender differentials. Occupation is an important socioeconomic indicator. The differences in prevalence of smoking were higher from one category of occupation to another. Prevalence among the people working in agriculture and as manual worker was high compare with domestic worker. A similar pattern of increasing in prevalence with age and decreasing with level of education was also observed with smokeless and current drinkers of alcohol. The habits of tobacco and alcohol use starts at early young age which contributes to the high risk of NCD at productive stage of life or as grown older with such habits.

The fruits and vegetable consumption and regular physical activity reduce the risk of non-communicable diseases. But, the study indicates high proportion of population taking inadequate amount of fruits and vegetables (less than five servings of fruits and vegetables per day). Low consumption across all the age groups, education and occupation by sex and residence was found high with marginal differences in between some of the groups. Besides that, more than half of the population was found in the category of doing

low physical activity. The differences in the pattern of low physical activity by age, sex, education, occupation and residence were also observed. Among the older age group (55-64), the people were doing less physical activity as compare with younger age. Female respondents were more in low physical activity compare with males across all the age groups. However, rural people were doing more physical work than urban, but similar pattern was observed across all age groups and sex. The physical activity by education was observed low among higher level education whereas it was comparatively high among lower level education people. Occupational difference in physical work activity was also observed across all the categories. The people whose occupation was agriculture or manual worker were doing more physical work compare with other occupational categories. Low physical activity was high among the domestic workers.

Hypertension is a major non-communicable disease risk factor especially related to cardiovascular disease. The increasing pattern of prevalence of hypertension (stage I & II) was recorded with increasing age group of people across all the subgroups of population (sex and urban-rural). It was prevalent in all the level of education with marginal differences with one another, which shows low impact of education on prevalence of hypertension. Hypertension was prevalent in all the occupational categories across residence and sex with some differences between the subgroups.

Overweight (obesity) is a major risk factor of Non-communicable diseases. High prevalence of overweight was recorded in all the age groups except the younger age (15-24). It was prevalent in both sexes, but higher in urban population compare with rural. Low prevalence of overweight was recorded among illiterate. It was low among the people working in agriculture or manual worker compare with other categories of occupation.

Overall, NCD risk factors were prevalent across all the socioeconomic and demographic categories of population in Andhra Pradesh.

The results generated through this IDSP-NCD survey would certainly focus on major health issues in bringing about change or initiate various programs related to control of non-communicable diseases.

CHAPTER 1

Introduction

1.1 BACKGROUND OF SURVEY

In response to a long felt need expressed by various expert committees, the Government of India through the Ministry of Health & Family Welfare initiated a decentralized, state based Integrated Disease Surveillance Project (IDSP) in the country with the assistance of the World Bank in the year 2004. The project envisaged detecting early warning signals of impending outbreaks; initiate an effective response in a timely manner. Unlike communicable diseases, most non-communicable diseases are latent type and they occur after a prolonged exposure to life style risk factors like smoking, raised blood sugar, raised blood pressure and hyper-cholestermia. Public health action would be primarily directed against preventive strategies for the disease and hence the priority was to monitor risk factors rather than non-communicable diseases themselves.

Periodic community based surveys covering representative adult population were planned under the IDSP to provide data on NCD risk factors at state level enabling states to develop strategies and activities to prevent and control the non-communicable diseases. It was taken up as a collaborative project of the Ministry of Health & Family Welfare, Govt. of India's National Institute of Communicable Diseases and the Indian Council of Medical Research with National Institute of Medical Statistics (NIMS) as the National Nodal Agency (NNA) and Regional Resource Centres (RRCs). The State Survey Agencies (SSAs) were identified based on their experience and knowledge about the local conditions. The division of non-communicable diseases at ICMR coordinated the overall activities and guided in the project development, implementation, monitoring and evaluation.

The NIMS provided the technical assistance at all stages of the survey including development of survey protocol, sampling methodology, survey questionnaire etc. with the approval of the National Technical Advisory Committee (NTAC). The survey was supervised and monitored by the RRCs for quality assurance. The RRCs

were identified in order to provide training to the field investigators, monitoring of data collection and technical support to the field agencies particularly for the anthropometrical and blood pressure measurements. All states were proposed to be covered in a phased manner. The first phase states included Andhra Pradesh, Kerala, Madhya Pradesh, Maharashtra, Mizoram, Tamil Nadu and Uttarakhand. The present treatise is the survey report of the state of Andhra Pradesh. Indian Institute of Health and Family Welfare, Hyderabad was the SSA in the state, while National Institute of Epidemiology, Chennai was the RRC.

1.2 OBJECTIVES

The overall objective of the NCD-risk factors survey was to improve the information available to the Government health services and care providers on a set of high-priority risk factors, with a view to improve on-the-ground responses to such risk factors. It also aimed to provide essential data to monitor progress of on going disease control programs and reallocate health resources more optimally. The specific objectives of the survey were to:

1. Assess the prevalence of NCD risk factors in different strata of population in the states;
2. Establish a baseline database of NCD risk factors needed to monitor trends in population health behavior and risk factors for chronic diseases over a period of time in the state; and
3. Provide evidence for evolving strategies and interventions for identified risk factors in the community to reduce the burden of Non-Communicable Diseases in the population

1.3 NON-COMMUNICABLE DISEASE (NCD) RISK FACTORS

A "risk factor" refers to any attribute, characteristic, or exposure of an individual, which increases the likelihood of developing a non-communicable disease. The major (modifiable)

behavioural risk factors identified in the World Health Report ² 2002 are tobacco use, harmful alcohol use, unhealthy diet (low fruit and vegetable consumption) and physical inactivity. On the other hand, the major biological risk factors identified are overweight and obesity, raised blood pressure, raised blood glucose and raised total cholesterol. These major behavioural and biological risk factors were included in non-communicable disease risk factors survey except raised blood sugar and total cholesterol, because they have the greatest impact on non-communicable disease mortality and morbidity, and modification is possible through effective prevention.

been identified as a risk factor in the development of a range of chronic diseases, including coronary heart disease, stroke and many forms of cancer. Research has indicated that the required intake of fruit for optimal health benefits is five daily servings of fruit and vegetable.

Lack of physical activity leads to obesity, dyslipidemia (lower high-density lipoprotein levels), insulin resistance, diabetes mellitus and high blood pressure levels. Physical inactivity is a well-established risk factor for coronary heart disease (CHD) and is associated with about a twofold increase in risk of CHD.

RISK FACTORS COMMON TO MAJOR NCD'S

Risk factor	Non-communicable Disease			
	CVD	Diabetes	Cancer	Respiratory
Smoking/tobacco	+	+	+	+
Alcohol	+		+	
Nutrition	+	+	+	+
Physical inactivity	+	+	+	+
Raised BP	+	+	+	
Raised blood sugar	+	+		
Obesity	+	+	+	+
Blood lipids*	+	+	+	

+ Corresponds to Risk Factor;

* Not being included in Phase I; CVD - Cardiovascular Disease

Tobacco use is a known or probable cause of about 25 diseases including heart disease; cancer, stroke, chronic obstructive pulmonary disease and digestive tract disease, as well as, has significant adverse effects on pregnancy. Smokeless tobacco use causes oral cancer in the lip, tongue, mouth, and throat areas and digestive system cancers. The relationship between alcohol consumption and health and social outcomes is complex and multi-dimensional. Alcohol consumption is linked to more than 60 disease conditions including liver cirrhosis, several cancers (liver, laryngeal, esophageal and oropharyngeal cancers), injuries and hemorrhagic strokes.

Consumption of fruits and vegetables reduces the risk of NCDs, like cancers and cardiovascular diseases. Dietary patterns that include higher intakes of fruits and vegetables are associated with several health benefits, including a decreased risk for some types of cancer. Low consumption of fruits and vegetables has

1.4 HEALTH PROFILE OF THE STATE

The state of Andhra Pradesh lies between 12° 41' and 22° longitude and 77° and 84° 40' latitude. It is the fifth largest state in India and it forms the major link between the north and the south of India. It is the biggest and most populous state in the south of India. The state is bounded by Madhya Pradesh and Orissa in the north, the Bay of Bengal in the east, Tamil Nadu and Karnataka in the south and Maharashtra in the west. It has an area of 275,045 sq. km. and a population of 81554 people (in thousand)³. There are 23 districts, 1127 blocks and 28123 villages in the state. The population density is 277 per sq. km. (as against the national average of 325). The population of the state has been growing with the decadal growth rate of 14.59% against 21.52% for the country. The key population and health indicators for Andhra Pradesh are presented in Table 1.1 and Table 1.2.

Table 1.1. Demographic and Socioeconomic profile of Andhra Pradesh as compared to India

S. No	Indicator	Andhra Pradesh	India
1	Total Population (in thousand)*	81554	1128521
2	Population Ratio (Urban /1000 Rural)*	376	385
3	Decadal Growth Rate*	14.59	21.52
4	Crude Birth Rate (Per 1000 Population)**	18.9	23.5
5	Crude Death Rate (Per 1000 Population)**	5.8	7.5
6	Life Expectancy at Birth**	62.7(M) 65.2(F)	62.3(M)63.9(F)
7	Total Fertility Rate***	2.0	2.9
8	Infant Mortality Rate (Per 1000 Live Births)**	56	57
9	Maternal Mortality Ratio (Per 100000 Live Births) †	195	301
10	Sex Ratio (Females/1000 Males)*	978	933
11	Mean Age at Marriage (Female)††	18.7	20.2
12	Population Below Poverty Line†††	15.8%	27.5%
13	Literacy Rate*	60.5%	64.8%

Source: National Health Profile 2007, Central Bureau of Health Intelligence⁴ (*Registrar General, India; **SRS Bulletin, October 2007; †Statistical Report, RGI 2004; ††Statistical Report RGI, 2005; RGI; PCA; †††Planning Commission of India).

Table 1.2. Health Infrastructure, Human Resource available and Health Expenditure

S. No	Indicator	Andhra Pradesh	India
1	Number of Allopathic Doctors with recognized medical qualifications and registered with State Medical Council*	48649	696747
2	Dental Surgeons Registered**	3726	72497
3	Number of Government Allopathic Doctors***	4487	76542
4	Average Population served/Doctor***	17988	-
5	Number of Registered AYUSH Doctors †	30049	725338
6	Total Number of Registered Nurses ††	189392	1509196
7	Number of Doctors at the PHCs †††	2202	22273
8	Total CHCs Specialists at CHCs †††	224	3979
9	Health Assistant (Male & Female) †††	3378	35330
10	Health Worker (Male & Female) †††	20067	215206

Source: National Health Profile 2007, Central Bureau of Health Intelligence, MOHFW

(*Medical Council of India; **Dental Council of India; ***Directorate of state health services; † Department of AYUSH, MOH&FW/GOI; ††Indian Nursing Council, Pharmacy Council of India; †††Bulletin on Rural Health Statistics in India, 2006 - Special Revised Edition, MOHFW)

1.5 SURVEY DESIGN AND IMPLEMENTATION

Sample Size

In order to achieve the aforesaid objectives, it was assumed that we should be able to estimate a parameter that has a level of 15% in a subgroup of population, with a relative precision of 30%, design effect as 1.25 and we would be able to achieve a response rate of 90%. Assuming that NCD risk factors are concentrated in 15-64 years for both males and females, the required sample size for each sex in 10-years age groups was estimated to be about 280. It is a known fact that the proportion of population in the 10 year age groups decrease with increase in age. In any population, the proportion of population in the age-group 55-64 is lowest

and varies in the range of 5-7 percent depending upon the fertility level (it is at the lower end, i.e., 5% in high fertility states, e.g. Uttar Pradesh, Madhya Pradesh, Bihar and Rajasthan, in the middle, i.e. 6% in moderate fertility states and at the upper end, i.e. 7% in low fertility states). Keeping such scenario of population composition in view and in order to have targeted 280 females and 280 males in age group 55-64, a sample of 5000 households was considered to be adequate for the survey.

Sample Design

A uniform sample design with equal allocation in urban and rural area was adopted in all states. In each state, the rural sample was selected in two stages: the selection of Primary Sampling Units (PSUs), which are

villages with probability proportional to population size (PPS) at the first stage, followed by the random selection of households within each PSUs at second stage using systematic random sampling. In urban areas, a three-stage procedure was followed. In the first stage, wards were selected with PPS sampling. In the second stage, one Census Enumeration Block (CEB) was randomly selected from each sample ward. In the final stage households were randomly selected within each CEB using the systematic random sampling procedure. From each selected PSU in rural area and from each selected Census Enumeration Block (CEB) in urban area, 50 households were selected. From each selected household, one individual was selected from those who fall in the 15-54 age range by using KISH method⁵ whereas all who fall in the age group 55-64 were included in the sample.

Sample Selection in Rural Areas

In rural area, the 2001 Census list of villages served as the sampling frame³. The list was stratified by a number of variables. The first level of stratification was geographic with villages classified into five contiguous regions. In each region, villages were further stratified by village size and the percentage of the population belonging to scheduled castes or scheduled tribes. The final level of stratification was implicit for all strata consisting of an ordering of villages within each stratum in ascending and descending order alternatively by the level of female literacy. From the list of villages so arranged, villages were selected systematically with probability proportional to the population of the village. Small villages with <75 households were linked with one or more adjoining villages to form PSUs. Villages with fewer than 5 households were excluded from the sampling frame.

In each selected sample PSU, a mapping and household listing was carried out prior to the data collection that provided the necessary frame for selecting households at the second stage. The household listing operation involved preparing up-to-date location map and layout sketch maps of each selected PSU, assigning numbers to structures, recording addresses or the location of these structures, identifying residential structures, and listing the names of the heads of all the households in residential structures in selected PSUs. The household listing operation was carried out by independent teams.

A complete listing of households was carried out in the villages with household up to 400 households. In case of villages with more than 400 households were

divided into at least three segments of 150-300 households as average size of each segment and two segments were selected for households listing using the systematic random sampling method. In each selected PSU, 50 households were selected from the household list using systematic random sampling.

Sample Selection in Urban Areas

The 2001 Census list of wards was used as the sampling frame. All wards were stratified by geographic regions, size of ward and percentage of SC/ST population. Female literacy was used for implicit stratification. A sample of wards was selected systematically with probability proportional of ward. One Census Enumeration Block (CEB), consisting of approximately 150-200 households, was selected from each selected ward using the PPS sampling method. The household listing operation was carried out in each selected census enumeration block similarly as in the village in rural area, which provided the necessary frame for selecting 50 households from the CEB.

Sample Weights

Appropriate sampling weights for households were used for urban and rural areas of the state. In urban sector it consisted of factors reflecting ward selection probabilities, Census enumeration block (CEB) selection probabilities within wards; and household selection probabilities within CEB; and household non-response adjustments. In rural sector, the element of weight consisted of factors reflecting probability of selection of PSU, household selection probability within the PSU, and household non-response adjustments.

From each selected household one member aged 15-54 was selected using the KISH method and all members aged 55-64 were selected. Since objective of the study was to obtain estimates for each age group (15-24 through 55-64) and by sex, post stratification was used for improvement of efficiency of the estimators. Post stratification weights for individuals are constructed using the state age distributions for both sexes of the urban and rural sector, which are available on the population level (Appendix-A)⁶.

Sample Implementation

During the survey, household information collected from a random sample of 4905 households covering 2431 households from rural and 2474 from urban areas. From these households, a total of 6270 respondents were contacted out of which 6249 completed the Step-1, and 6218 completed the Step-2 survey. The

overall individual non-response rate for the survey was less than 1% (Table 1.3)

Against the target sample size of 280, there is low turnout in certain age groups and high turn out in other (it may be seen in the subsequent table 2.2). It might be due to either misreporting of age or replacement of the individual who was selected but not available at the

to the head of the household. The residential status (whether present in the household or temporary away from household) was gathered. The above information were used to identify the eligible individual for the survey in the age group 15-64 years, for administering individual questionnaire. The Household Questionnaire also collected information on religion, ownership of a

Table 1.3 Sample coverage and response rate of household, step-1 and step-2 individual response rate by place of residence, Andhra Pradesh, 2007- 08

Response	Residence		
	Urban	Rural	Combined
Households interview			
Households contacted	2500	2500	5000
Households interviewed	2474	2431	4905
Households response rate (%)	99.0	97.2	98.1
Eligible Participants Step-1			
Individual contacted	2776	3494	6270
Individual interviewed	2764	3485	6249
Response rate (%)	99.6	99.7	99.7
Eligible Participants Step-2			
Step-2 completed	2750	3468	6218
Overall individual response rate (%)	99.0	99.3	99.2

time of interview affecting the use of Kish method to give the required sample size. In fact, the Kish method was used in each selected household to select one respondent amongst those who were aged 15-54. It was done by the field investigator after listing of members of the household and arranging them according to age, sex and then selecting one respondent for the interview. There is possibility that some respondents in the age group 15-54 particularly males were not available at home during the survey (10AM to 5 PM) and thereby might have been replaced by those household members who were present at the time of survey.

1.6 SURVEY INSTRUMENTS

The survey used two types of questionnaire, the Household Questionnaire and the Individual Questionnaire (Appendix-B). The overall content and format of the questionnaires were determined through a series of workshops and meetings held in 2006-07. The questionnaires for each state were bilingual with questions in both the English and principal language of the state which was Telegu in the present case. It first listed all usual residents age 12 years and above, in each sample household. For each listed members, survey collected basic information on age, sex and relationship

house, type of house with number of rooms, main source of drinking water, type of toilet facility, main source of lighting, types of cooking fuel, type of oil/cooking medium, ownership of agricultural land, ownership of livestock and possession of durable goods.

The Individual Questionnaire included questions seeking information from all the selected individuals (men and women) in the age group 15-64. The Individual Questionnaire covered information on demographic, behavioural and physical measurements under Step-1 and Step-2 with a number of sections into them. The first section of Step-1 included questions regarding the demographic information of individual, i.e., age, sex, marital status, education, and occupation. The behavioural information section included questions on tobacco use, alcohol consumption, diet, and physical activity, history of raised blood pressure and history of diabetes.

Tobacco Use (Smoking & Smokeless): Questionnaire was used to elicit information on current and past use of tobacco (smoking & smokeless), age when used tobacco for first time, past history of tobacco use, and age when stopped using tobacco.

Alcohol use: Questionnaire collected information on

whether the individual was currently using alcohol, use of alcohol in past 12 months, frequency of drinks in past 12 months, average number of drinks consumed in one day, alcohol consumed within past 30 days, number of standard alcoholic drink consumed per day in past 7 days, past history of alcohol consumption, and age when started consuming alcohol regularly.

The contents and format of these questionnaires were though largely governed by the WHO STEPS guidelines but they were finalized through a series of consultative meetings held at the Indian Council of Medical Research.

Diet: Questions were asked to collect information on number of days in a week when fruits were consumed, number of serving of fruits consumed in a day, number of days in a week when vegetables were consumed, number of servings of vegetables consumed in a day, frequency of consumption of cheese and butter, fried local food, red meat, eggs, chicken, fish, aerated soda, sweetened drinks, pizza/burger/French fries, cakes/pastries or other bakery items, chips/*namkeen*.

Physical Activity: Questions were asked about the intensity of physical activity in the daily work, frequency of doing physical activity of varying intensity, time spent in doing physical activity of varying intensity per day, mode of travel to and from places, time spent walking or bicycling, type of vigorous/moderate intensity sports for recreation being practiced, frequency of doing such vigorous/moderate intensity sports in a week, time spent doing vigorous/moderate intensity sports per day, practice of yoga, frequency of practicing yoga, duration of time spent per day in yoga, time spent sitting or reclining etc.

History of Raised Blood Pressure: Questions were asked on history of hypertension, medicines prescribed by a doctor and the advice given regarding diet, weight lose, smoking and nature of physical activity undertaken.

History of Diabetes: Questions covering history of diabetes, medicines prescribed by a doctor and advice given regarding diet, weight lose, smoking and physical activities were asked.

Individual questionnaire included several biomarker measurements in Step-2. The height of the eligible individual participant was taken in centimeter by using a portable height measuring board and also measured weight in kilogram using a portable electronic weighting scale. Waist circumference (not measured for pregnant women) was taken two times to provide additional

information on overweight and obesity. Constant tension tape (Figure finder tape) measure was used for waist circumference measurements. The measurement is taken without clothing, that is, directly over the skin or over light clothing. The privacy area was maintained for this measurement.

Blood pressure of the individual participants was taken three times using automated blood pressure measuring instrument (OMRON) and pulse rate was also measured three times using an automated blood pressure device.

1.7 TRAINING

In order to maintain uniform survey procedure across the country, a manual dealing with various aspects of the survey were prepared by NIMS, ICMR. There are five sections: (1) Project Protocol, (2) Survey Methodology, (3) Coordinator's Guide, (4) Trainers Guide and (5) Interviewer's Guide. The Interviewer's Guide consists of guidelines to the interviewers regarding interviewing procedure, field procedures and method on asking each question and recording answers. The Coordinator's Guide contains a detail description of the role and responsibilities of the state coordinators. The Trainer's Guide include training guidelines for the training of the field staff including survey methodology, survey instruments, mapping and list of households, preparation and collection of data.

The representatives of State Survey Agencies (SSAs) and Regional Resource Centres (RRCs) were trained in the Training of Trainers workshop and Data Entry & Management Workshop organized by NIMS at the beginning of the data collection (18-20 July 2007). The purpose of the former workshop was to explain the objective of the NCD Risk Factors Survey and ensure uniform application of survey material to collect good quality of data. The survey documents such as training manuals, survey instruments, list of selected rural and urban PSUs etc. was provided to them for conducting the survey. The equipments required for survey was procured centrally by ICMR and distributed to the SSAs and RRCs. The personnel trained in these workshops subsequently trained the field staff in their respective states.

Training of Field Staff

As mentioned, the field staff recruited for the survey in Andhra Pradesh was trained by IIHFW, Hyderabad and the officials of NIMS, New Delhi and NIE, Chennai, supervised the training process. The training

was conducted from 1st - 6th October, 2007 at IIFHW, Hyderabad. The training consisted of lectures, classroom training, demonstration, practice interviews and field based training. A total of 30 participants were trained, of these 30 trainees 25 were part of the 5 survey teams and 5 were the supervisors for the survey teams. It was ensured that each survey team comprised of one male and one female member.

Each trainee was given a training kit at the beginning of training, the training kit comprised of an interviewers guide, household and individual schedules, consent form, IEC message, set of show cards (e.g. diet chart, alcohol chart) and reference forms (e.g. Kish table, table of random numbers), flow chart of activities in field, identity card and supporting letters from Government mentioning purpose of visit. A field visit to village Patanchervu, Medak district was also arranged as part of practical training of investigators in field activities and procedure for conducting a survey and as part of pre-test. After the completion of training, letters were issued through Directorate of Health Services to DHOs, Municipal corporations and Municipalities across the state, and Deputy Director of Health Services requesting their cooperation in smooth conduction of the survey.

Data Entry Training

Data entry software in Epi-Info with its manual was developed by the NIMS, ICMR. A two-day data entry workshop cum hands on training was organized by NIMS, ICMR during 10-11 December 2007 for the statisticians and data entry personnel of the state survey agencies (SSAs). The purpose of the workshop was to familiarize the participants with the software. Emphasis was made on double data entry in order to ensure high accuracy in data entry and to minimize data entry errors. All the participants were provided with the Data Entry Software and its Manual.

1.8 DATA PROCESSING AND ANALYSIS

Following the data entry by the state survey agency (SSA), the validity and consistency check was carried out by the NIMS, ICMR for final analysis. Analysis plan in terms of dummy tables was finalized in consultation with ICMR Review Group.

Prevalence of current smokers, current daily

smokers and past daily smokers was calculated among the respondents by sex and place of residence. Those who smoke tobacco daily, the mean number of tobacco products (*bidis, cigarettes, pipes, cigars, etc.*) used daily was calculated taking denominator as all daily smokers. Though the age of initiation of smoking was collected from all daily smokers in completed years but for the past smokers it was calculated by imputation because it was not recorded. Finally average age of initiation of smoking was calculated in two age groups of smokers, 15-34 years and 35-64 years. The same procedure was followed for the calculation of average age of initiation of smokeless tobacco. Prevalence of alcohol consumption was calculated for last twelve months, last thirty days and last seven days and presented as percentage. The mean age of initiation of alcohol consumption was also calculated. Mean number of servings fruit, vegetables, and combined (fruit and vegetables) consumed per day was computed.

Mean physical activity per day was computed by combining all types of physical activity (vigorous, moderate-intensity, travel and recreational) using METs (Metabolic Equivalent) score. Prevalence of reported cases of blood pressure and diabetes were also calculated. Measurement of height, weight and waist circumference of individual respondent was used to compute BMI (body mass index) and central obesity.

1.9 QUALITY CONTROL MEASURES

A uniform project protocol, survey methodology, training manuals, survey instruments and data-management modules were developed and adopted across all the states. It was executed by the Indian Institute of Health and Family Welfare, Hyderabad (SSA) and was monitored by National Institute of Epidemiology, Chennai (RRC). The overall coordination and supervision of the survey was done by the Indian Council of Medical Research and the National Institute Medical Statistics, New Delhi. In addition, an independent check by collecting data in randomly drawn sample of 10% of PSUs was carried out by the RRC. High concordance was recorded between the survey by SSA and independent checked by RRC on some key indicators like smoking, alcohol consumption and physical activity with an overlapping of 95% confidence interval. Various activities hitherto were to maintain the highest level of the quality of data.



CHAPTER 2

Background Characteristics of the Households and Respondents

This chapter presents the demographic and socio-economic characteristics of the sample households and the respondents from these households in the survey population of Andhra Pradesh. It also describes facilities in the households. The overall prevalence of various risk factors is presented for the ages 15-64 years.

2.1 HOUSEHOLD CHARACTERISTICS

Table 2.1 provides the percentage distribution of households in rural and urban areas by various characteristics of the surveyed households. Majority of the households were Hindu (82%) followed by Christian (11%) and Muslim (8%). Seventy five percent of sample households had piped drinking water supply followed by 16% from hand pump, 8% from well and a small fraction (0.2%) from surface. It is found that 94% of urban households and 68% of rural households had piped

drinking water supply. Regarding the sanitation facility, 47% of the households had flush toilets and 53% had pit toilets. The urban households were more likely to have access to the flush toilets (89%) as compared to rural households (32%) in Andhra Pradesh.

Over ninety percent (93%) households use electricity as main source of lighting which was slightly higher in urban area (98%) than that in the rural area (91%). In the state, 54% households have *pucca* house, 27% households have *semi-pucca* house and 19% households have *kaccha* house. High percentage of households in the urban area (70%) compared to the rural area (49%) have *pucca* houses. Several types of fuel were used for cooking in Andhra Pradesh, with wood as the most common type (81%) in the rural areas L.P.G. was most common (72%) in urban area. In the state as whole, 65% households used wood followed by LPG (31%) and kerosene (2%).

Table 2.1 Percentage distribution of households in rural and urban area according to the background characteristics, Andhra Pradesh, 2007- 08

Characteristics	Residence		Combined
	Urban	Rural	
Religion of household head			
Hindu	81.6	81.7	81.7
Muslim	12.2	5.8	7.5
Christian	6.1	12.5	10.8
Total	100.0	100.0	100.0
Source of drinking water			
Piped	93.9	68.2	74.9
Hand pump	4.9	19.4	15.6
Well water	0.3	10.8	8.1
Surface water	0.0	0.2	0.2
Other	0.9	1.4	1.2
Total	100.0	100.0	100.0
Sanitation facility			
Flush toilet	88.5	32.1	46.8
Pit toilet	11.5	67.8	53.2
Total	100.0	100.0	100.0
Main source of lighting			
Electricity	98.7	91.3	93.2
Kerosene	1.1	8.5	6.6

Gas/Oil	0.2	0.2	0.2
Total	100.0	100.0	100.0
Type of house			
<i>Pucca</i>	69.7	48.8	54.2
<i>Semi-Pucca</i>	25.9	27.5	27.1
<i>Kachha</i>	4.5	23.7	18.7
Total	100.0	100.0	100.0
Cooking fuel			
LPG	72.0	17.1	31.3
Wood	21.0	80.8	65.3
Kerosene	4.8	0.4	1.6
Others	2.1	1.7	1.8
Total	100.0	100.0	100.0
Separate kitchen room			
Yes	60.3	37.8	43.7
No	39.7	62.2	56.3
Total	100.0	100.0	100.0
Agriculture land			
Own agriculture land (%)	6.7	36.8	28.9
Number	2474	2431	4905

Seven in every ten households in Andhra Pradesh had no own agricultural land. Three out of every five households in rural area had no own agricultural land compared to 93% in urban area. The proportion of households having separate kitchen was 44%. This percentage was 60 in urban and 38 in rural area.

2.2 AGE AND SEX COMPOSITION

A total of 6218 with 2719 males and 3499 females from urban and rural areas were contacted in the survey. They are presented in 10 years age groups 15-24, 25-34, 35-44, 45-54 and 55-64. The distribution of the number of respondents across the five age groups is depicted for both males and females as well as for both sex together, separately for urban and rural areas and combined in Table 2.2. It may be seen that the number

of respondents was least, i.e. 1171, in the age group 45-54 and maximum, i.e., 1560 in age group 25-34 years.

2.3 EDUCATION LEVEL

Table 2.3 presents the percentage of the respondents according to their literacy levels by sex and place of residence. Forty five percent of the total respondents were illiterate whereas 22% were with primary or middle, 24% were secondary or in higher secondary, while 9% were educated up to college and above. In the sample, 56% females and 34% males were illiterate. The proportion of illiterate respondents among rural females was as about the twice as high than that among urban females. A higher percentage of males than of females had completed almost each level of schooling. Education levels were much higher for urban respondents than for rural respondents.

Table 2.2 Age and Gender-wise distribution of respondents by type of residence (unweighted), Andhra Pradesh, 2007- 08

Age	Residence						Combined		
	Urban			Rural					
	Male	Female	Total	Male	Female	Total	Male	Female	Total
15 - 24	227	302	529	236	350	586	463	652	1115
25 - 34	304	439	743	339	478	817	643	917	1560
35 - 44	291	301	592	302	384	686	593	685	1278
45 - 54	238	224	462	315	394	709	553	618	1171
55 - 64	197	227	424	270	400	670	467	627	1094
15 - 64	1257	1493	2750	1462	2006	3468	2719	3499	6218

2.4 MARITAL STATUS

The second panel of Table 2.3 shows the percentage of respondents according to the three marital categories by sex and place of residence. Three quarter of the respondents, both male and female were currently married, 9% were widow/divorced or separated and 20% were never married. The proportion of respondents who were currently married did not vary much by urban rural residents.

2.5 OCCUPATION

Table 2.3 provides information on the current occupation of the respondents. In the sample about half of the females (50%) are engaged in domestic works. It was closely followed by individuals involved in agricultural works, which were about 32% in case of males and 25% in case of females. About 20% individuals were engaged in manual work and 9% were engaged in executive/business work.

Table 2.3 Percentage of respondents according to background characteristics, gender and place of residence, Andhra Pradesh, 2007- 08

Characteristic	Residence						Combined		
	Urban			Rural			Male	Female	Total
	Male	Female	Total	Male	Female	Total			
Education									
Illiterate	16.8	35.9	26.1	40.6	64.0	52.3	33.7	56.1	44.8
Primary	6.1	8.2	7.2	10.6	9.2	9.9	9.3	8.9	9.1
Middle	14.3	14	14.2	13.7	11.1	12.4	13.9	11.9	12.9
Secondary	21.7	19.6	20.6	18.9	10.2	14.6	19.7	12.8	16.3
Higher Secondary	16.3	10.4	13.4	8.6	3.4	6.0	10.8	5.4	8.2
College & above	24.8	11.9	18.5	7.6	2.1	4.8	12.6	4.9	8.7
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Marital status									
Never married	30.9	14.6	23	26.3	10.9	18.6	27.6	11.9	19.8
Married	67.6	75.8	71.6	70.1	73.3	71.7	69.4	74.0	71.7
Widowed/Divorced/ Separated	1.5	9.6	5.4	3.6	15.8	9.7	3.0	14.1	8.5
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Occupation									
Executive/Business	29.4	6.6	18.3	8.1	3.4	5.7	14.2	4.3	9.3
Agriculture	5.2	4.3	4.8	43.3	33.3	38.4	32.3	25.2	28.8
Domestic work	0.4	67.5	33.1	0.2	42.8	21.4	0.3	49.7	24.8
Services/Sales	11.6	3.8	7.8	5.7	1.8	3.7	7.4	2.3	4.9
Manual worker	31.7	7.8	20.0	27.8	12.6	20.2	28.9	11.3	20.1
Other	21.7	10.0	16.0	14.9	6.1	10.6	16.9	7.2	12.1
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Number	1257	1493	2750	1462	2006	3468	2719	3499	6218



CHAPTER 3

Behavioural Risk Factors

This chapter presents the prevalence of certain behavioural risk factors for the non-communicable diseases in the survey population. The survey questionnaire asked questions about certain life style of respondents which could be considered as the behavioural risk factors for non-communicable diseases.

3.1 TOBACCO SMOKING AND CHEWING

As per the WHO STEPS guidelines to measure the prevalence of smoking habit among the respondents, the smokers are categorized as *current smokers*, *current daily smokers*, *past daily smokers* and those who have never smoked in lifetime are classified as *non-smokers*.

Table 3.1.1 presents the percentage of respondents, both males and females as well as combined in various categories of smokers and non-smokers according to the place of residence (rural or urban). About 18% of the respondents in the survey were current smokers. They were mostly current daily smokers. The prevalence of smoking was more among men (32%) than among women (4%). By place of residence, 12% of urban

respondents and 19% of rural respondents were current daily smokers. Only 3% of respondents were past daily smokers and 79% of respondents had never smoked tobacco.

Table 3.1.2 presents the mean number of smoked among those respondents who were current daily smokers of any form of tobacco. Among those who were current daily smoker of different product of tobacco, the mean number of smoking per day was 5 in case of *bidis* and 3 for manufactured cigarettes. The bidi and manufactured cigarettes smokers were mostly males. Among those who smoke bidi, the frequency of smoking was higher among rural respondents (5) as compared to urban respondents (3). In case of manufactured cigarettes however, the mean number of smoking was high among urban respondents (5) as compared to rural respondents (3).

Table 3.1.3 presents the mean age of initiation, age at stopped smoking and the percentage of non-smoker respondents exposed to tobacco smoke by sex and the place of residence.

The mean age for initiation of smoking among young respondents aged 15-34 years was 19 years and

Table 3.1.1 Percentage of respondents classified by smoking status across sex and place of residence, Andhra Pradesh, 2007- 08

Residence/ Sex	Smoking Status							
	Current smokers		Current daily smokers		Past daily smokers		Never smoked	
	P (%)	95% CI	P (%)	95% CI	P (%)	95% CI	P (%)	95% CI
Urban								
Male	23.9	(20.2,28.2)	23.0	(19.2,27.3)	6.5	(5.1,8.4)	69.5	(65.2,73.5)
Female	0.8	(0.5,1.4)	0.8	(0.5,1.4)	0.3	(0.1,0.8)	98.9	(98.1,99.3)
Total	12.6	(10.7,14.8)	12.2	(10.3,14.4)	3.5	(2.7,4.5)	83.8	(81.5,85.9)
Rural								
Male	34.5	(29.9,39.5)	33.9	(29.2,38.9)	4.3	(3.1,6.0)	61.1	(56.0,66.0)
Female	5.2	(3.3,8.3)	4.9	(2.9,8.0)	0.7	(0.4,1.2)	94.1	(90.9,96.2)
Total	19.9	(17.5,22.6)	19.4	(16.9,22.2)	2.5	(1.9,3.4)	77.6	(75.0,79.9)
Combined								
Male	31.5	(28.0,35.2)	30.7	(27.2,34.5)	5.0	(4.0,6.2)	63.6	(59.8,67.2)
Female	4.0	(2.6,6.2)	3.7	(2.3,5.9)	0.6	(0.3,1.0)	95.4	(93.1,97.0)
Total	17.9	(16.0,19.8)	17.4	(15.5,19.4)	2.8	(2.3,3.5)	79.3	(77.4,81.1)

Table 3.1.2 Mean number of tobacco products smoked (per day) by daily smokers according to sex and place of residence, Andhra Pradesh, 2007- 08

Residence/ Sex	Type of smoking			
	Bidi		Manufactured Cigarettes	
	Mean	95% CI	Mean	95% CI
Urban				
Male	3.6	(2.4, 4.7)	5.3	(4.5, 6.2)
Female	*		0.8	(0.0, 2.2)
Total	3.4	(2.3, 4.6)	5.2	(4.4, 6)
Rural				
Male	5.4	(4, 7)	3.1	(2.4, 4.0)
Female	0.6	(0.0, 1.7)	0.0	(0, 0)
Total	4.8	(3.4, 6.2)	2.8	(2, 3.4)
Combined				
Male	5.0	(4, 6)	3.6	(3, 4)
Female	0.5	(0.0, 1.5)	0.1	(0.0, 0.1)
Total	4.5	(3.4, 5.7)	3.2	(2.7, 3.8)

*no observation

Table 3.1.3 Mean age of initiation, age at stopped smoking and percentage of respondents (non-smokers) exposed to tobacco smoke by sex and place of residence, Andhra Pradesh, 2007- 08

Residence/ Sex	Smokers						Non-smokers	
	Age of Initiation (15-34 years)		Age of Initiation (35-64 years)		Age at Stopped		Exposed to tobacco smoke at Home or Work	
	Mean	95% CI	Mean	95% CI	Mean	95% CI	P (%)	95% CI
Urban								
Male	19	(18.5, 20.5)	20	(19.0.5, 20.5)	32	(29, 35)	28	(23.7, 33.7)
Female	*		25	(20.5, 30.5)	37	(30, 45)	12	(9.4, 16.9)
Total	19	(18.5, 20.5)	20	(19.5, 20.5)	32	(29, 35)	19	(16.5, 23.2)
Rural								
Male	19	(17.5, 20.5)	20	(19.0, 20.5)	32	(28, 37)	20	(15.0, 26.1)
Female	14	(10.5, 25.5)	16	(14.5, 17.5)	32	(22, 41)	19	(14.6, 24.9)
Total	19	(17.5, 20.5)	20	(18.5, 20.5)	32	(28, 37)	20	(15.3, 24.9)
Combined								
Male	19	(18.5, 20.5)	20	(19.0, 20.5)	32	(29, 35)	23	(19.1, 27.3)
Female	14	(10.5, 25.5)	16	(15.5, 18.5)	32	(24, 41)	17	(13.9, 21.4)
Total	19	(18.5, 20.5)	20	(19.5, 20.5)	32	(29, 35)	20	(16.5, 23.4)

*no observation

among all respondents aged 35-64 years was 20 years. The mean age of cessation of smoking for all those who stopped smoking was 32 years. The age of initiation of smoking in the age group 35-64 years was 20 years for rural male and 16 years for rural female respondents whereas among the urban smokers the mean age of initiation for male was 20 years and for female 25 years. The mean age of cessation of smoking for urban females was 37 years against 32 years for rural females.

About 20% respondents of those who never smoked were exposed to tobacco smoke at home or work place.

It was 23% in case of men against 17% in case of women. The urban and rural differentials were low with respect to exposure to tobacco smoke at home or work place.

Table 3.1.4 provides percentage of smokeless tobacco users by sex and place of residence of the respondents.

One-tenth of the respondents were current user of smokeless tobacco with 14% among men and 5% among women. A small percentage of men and women (1%) were found to be past daily users. The prevalence of smokeless tobacco use was similar in rural and urban

Table 3.1.4 Percentage of smokeless tobacco users by sex and place of residence, Andhra Pradesh, 2007- 08

Residence/ Sex	Smokeless tobacco user							
	Current user		Current daily user		Past daily user		Never used	
	P (%)	95% CI	P (%)	95% CI	P (%)	95% CI	P (%)	95% CI
Urban								
Male	15.2	(11.7, 19.4)	14.3	(10.9, 18.4)	2.2	(1.3, 3.2)	82.5	(78.6, 86.6)
Female	3.0	(1.8, 5.0)	3.0	(1.8, 5.0)	0.2	(0.0, 0.5)	96.7	(95.1, 98.3)
Total	9.2	(7.1, 11.9)	8.8	(6.7, 11.4)	1.3	(0.8, 1.7)	89.5	(86.9, 92.0)
Rural								
Male	13.0	(9.2, 18.0)	12.2	(8.4, 17.3)	1.6	(0.5, 2.7)	85.4	(80.7, 90.1)
Female	5.1	(2.9, 8.9)	5.1	(2.9, 8.8)	0.2	(0.0, 0.5)	94.6	(91.7, 97.5)
Total	9.1	(7.0, 11.7)	8.6	(6.5, 11.3)	0.9	(0.4, 1.0)	90.0	(87.5, 92.4)
Combined								
Male	13.6	(10.7, 17.2)	12.8	(9.9, 16.4)	1.8	(0.9, 2.6)	84.6	(81.1, 88.0)
Female	4.5	(2.9, 7.1)	4.5	(2.8, 7.1)	0.5	(0.0, 0.5)	95.2	(93.1, 97.3)
Total	9.1	(7.5, 11.1)	8.7	(7.0, 10.6)	1.0	(0.6, 1.4)	89.8	(88.0, 91.7)

areas (9%). More men (12% in rural area 14% in urban area) than women (5% in rural area and 3% in urban area) were current daily users of smokeless tobacco.

The mean number of consumptions per day of various smokeless tobacco products such as tobacco chewing, pan with tobacco, snuff by mouth, snuff by nose and others are provided in Table 3.1.5.

The mean number of times chewing tobacco, *Pan* with tobacco and snuff by mouth/nose per day in Andhra Pradesh was less than 1. There was no urban-rural and male-female differential in frequency of chewing tobacco. For those who used snuff by mouth the average daily frequency of such consumption were around 2.

The mean age of initiation and age at stopped using

of smokeless tobacco use by sex and place of residence of respondents is provided in Table 3.1.6. The mean age of initiation of smokeless tobacco was 20 years among aged 15-34 years. While there was no sex differential in mean age of initiation of smokeless tobacco use in urban area, rural male appears to initiate smokeless tobacco slightly earlier (20 years) than rural female (23 years). For respondents aged 35-64 years, the mean age of initiation of smokeless tobacco use was 25 years (25 years for males and 20 years for females). In urban population aged 35-64, the mean age of initiation of smokeless tobacco was 24 years (24 for males and 25 for females) and among rural respondents it was 25 years (30 for the males and 20 for the females). The mean age of quitting smokeless

Table 3.1.5 Mean frequency of smokeless tobacco use (per day) by daily smokeless tobacco users according to sex and place of residence, Andhra Pradesh, 2007- 08

Residence/ Sex	Type of smokeless tobacco								Others	
	Chewing tobacco		<i>Pan</i> with tobacco		Snuff by mouth*		Snuff by nose			
	Mean	95% CI	Mean	95% CI	Mean	95% CI	Mean	95% CI	Mean	95% CI
Urban										
Male	0.5	(0.2, 0.7)	1.6	(1.0, 2.1)	2.1	(1.5, 2.8)	0.6	(0.0, 1.2)	0.4	(0.0, 0.8)
Female	0.7	(0.0, 1.3)	1.6	(0.9, 2.3)	1.6	(1.0, 2.4)	0.6	(0.3, 1.6)	0.1	(0.0, 0.2)
Total	0.5	(0.2, 0.7)	1.6	(1.1, 2.0)	2.0	(1.4, 2.7)	0.6	(0.0, 1.1)	0.4	(0.0, 0.7)
Rural										
Male	0.9	(0.5, 1.3)	0.6	(0.3, 1.0)	2.0	(1.3, 2.6)	0.6	(0.7, 1.9)	0.3	(0.0, 0.5)
Female	1.0	(0.6, 1.4)	1.4	(0.5, 2.2)	1.0	(0.4, 1.3)	0.7	(0.4, 1.8)	0.0	(0.0, 0.7)
Total	0.9	(0.6, 1.2)	0.9	(0.5, 1.2)	1.7	(1.1, 2.2)	0.2	(1.4, 0.6)	0.2	(0.0, 0.4)
Combined										
Male	0.8	(0.5, 1.0)	1.0	(0.6, 1.3)	2.0	(1.6, 2.5)	0.0	(0.0, 0.2)	0.3	(0.1, 0.6)
Female	0.9	(0.6, 1.3)	1.4	(0.7, 2.1)	1.0	(0.7, 1.4)	0.6	(0.3, 1.5)	0.0	(0.0, 0.0)
Total	0.8	(0.6, 1.0)	1.0	(0.8, 1.4)	1.8	(1.4, 2.2)	0.2	(0.0, 0.5)	0.2	(0.0, 0.4)

*Tooth powder or tooth paste prepared using tobacco

Table 3.1.6 Mean age of initiation, age at stopped smokeless tobacco use by daily smokeless tobacco user according to sex and place of residence, Andhra Pradesh, 2007- 08

Residence/ Sex	Smokeless tobacco users					
	Age of Initiation (15-34 years)		Age of Initiation (35-64 years)		Age at Stopped	
	Mean	95% CI	Mean	95% CI	Mean	95% CI
Urban						
Male	20	(18.5, 21.5)	24	(22.5, 25.5)	28	(23.8, 32.1)
Female	20	(12.5, 25.5)	25	(20.5, 30.5)	26	(20.8, 32.0)
Total	20	(19.5, 21.5)	24	(22.5, 25.5)	28	(24.0, 32.0)
Rural						
Male	20	(18.5, 22.5)	30	(25.5, 33.5)	25	(21.0, 30.30)
Female	23	(10.5, 28.5)	20	(19.5, 25.5)	37	(24.3, 52.0)
Total	21	(20.5, 22.5)	25	(20.5, 28.5)	27	(22.5, 31.2)
Combined						
Male	20	(19.5, 21.5)	25	(24.5, 27.5)	26	(23.2, 29.4)
Female	23	(19.5,27.5)	20	(19.5, 25.5)	34	(26.0, 42.5)
Total	20	(19.5, 21.5)	25	(22.5, 25.5)	27	(24.1, 30.1)

tobacco use for those who did so was 27 years; it was 26 years in case of men and 34 years in case of women. The mean age of quitting smokeless tobacco use for urban respondents was 28 years against 27 years for rural respondents. There was no much difference in the age of quitting smokeless tobacco between male and female urban respondents (28 years for males against 26 years for females), but there was marked difference in rural male and female respondents (25 years for males against 37 years for females).

Table 3.1.7 presents the percentage of smokers and smokeless tobacco users by sex and the place of residence. It shows that 24% of respondents were either smoking or use smokeless tobacco whereas 2% of the

respondents were using both forms of tobacco, i.e. smoking and also smokeless tobacco. The use of either smoking or smokeless tobacco was 19% for urban as compared to 26% for rural. The use of both the forms of tobacco (smoking as well as smokeless tobacco) was 2% for urban and 3% for rural respondents. The percentage of using either smoking or smokeless tobacco among men was high (39%) as compared to women (8%).

Some key observations on tobacco use was that one-third of male population smoked tobacco daily whereas smoking among females was low. Overall 9% of the adult population used smokeless tobacco with 13% of men and 5% of women used smokeless tobacco. Twenty four percent of adult population in Andhra

Table 3.1.7 Percentage of tobacco users by sex and place of residence, Andhra Pradesh, 2007- 08

Residence/ Sex	Tobacco Use							
	Smokeless tobacco users only		Smokers only		Both (Smoking and smokeless)		Either (Smoking or smokeless)	
	P (%)	95% CI	P(%)	95% CI	P (%)	95% CI	P (%)	95% CI
Urban								
Male	10.5	(7.9, 13.8)	19.2	(16.2, 22.7)	3.8	(2.3, 6.2)	33.5	(29.1, 38.2)
Female	3.0	(1.8, 5.0)	0.8	(0.5, 1.3)	0.0	(0.0, 0.2)	3.8	(2.5, 5.7)
Total	6.8	(5.1, 9.0)	10.2	(8.7, 12.0)	2.0	(1.2, 3.2)	19.0	(16.5, 21.8)
Rural								
Male	7.5	(4.9, 11.3)	29.2	(24.2, 34.8)	4.7	(3.2, 6.8)	41.4	(37.2, 45.7)
Female	4.8	(2.7, 8.4)	4.6	(2.7, 7.8)	0.2	(0.1, 0.7)	9.7	(7.0, 13.4)
Total	6.2	(4.4, 8.5)	17.0	(14.6, 19.7)	2.5	(1.7, 3.5)	25.6	(23.2, 28.1)
Combined								
Male	8.4	(6.3, 11.0)	26.3	(22.7, 30.3)	4.4	(3.2, 6.0)	39.1	(35.8, 42.5)
Female	4.3	(2.7, 6.8)	3.5	(2.1, 5.8)	0.2	(0.1, 0.5)	8.0	(6.0, 10.7)
Total	6.4	(5.0, 8.1)	15.0	(13.3, 17.0)	2.3	(1.7, 3.1)	23.7	(21.8, 25.7)

Pradesh used tobacco in any form (i.e. smoking or smokeless). The mean age of initiation of tobacco use among young age (15-34 years) people was 19 years for male smokers, and 20 years for male smokeless tobacco users. These findings emphasize the need of implementing the tobacco control programme for prevention of NCD.

3.2 ALCOHOL CONSUMPTION

Table 3.2.1 presents the percentage of respondents who consumed alcohol in past 30 days and 12 months by sex and place of residence.

About 14% respondents had consumed alcohol in past 30 days and 20% consumed in past 12 months. Only 3% respondents were past drinker. Twenty seven percent men consumed alcohol in past 30 days and about 37% men did so in past 12 months as compared to very less

among women (3%). Percentage of lifetime abstainer to alcohol was high for rural (79%) as compared to urban respondents (74%). Urban men were more likely to consume alcohol (29% in past 30 days and 36% in past 12 months) than rural men (26% in past 30 days and 36% in past 12 months).

Table 3.2.2 presents the percentage of those who consumed alcohol according to frequency of consuming alcohol in past 12 months, mean number of standard drinks consumed on a drinking day, frequency of consuming alcohol in past one week and the mean number of standard drinks per day.

Of the current drinkers, 33% in urban area, 27% in rural area and 29% in the combined sample consumed alcohol on less than one occasion in a month. Further, 36% of urban and 28% of rural such respondents consumed alcohol 1-3 days per month in past one year. Twenty

Table 3.2.1 Percentage of Alcohol consumption by sex and place of residence, Andhra Pradesh, 2007- 08

Residence/ Sex	Alcohol Consumption							
	Consumed alcohol (Last 30 days)		Consumed alcohol (Last 12 months)		Past drinker		Life time abstainer	
	P (%)	95% CI	P (%)	95% CI	P (%)	95% CI	P (%)	95% CI
Urban								
Male	28.7	(24.2, 33.6)	36.2	(34.3, 45.5)	8.9	(6.3, 12.5)	54.9	(49.2, 60.5)
Female	1.7	(0.9, 3.3)	3.5	(2.2, 5.8)	0.5	(0.2, 1.3)	96.0	(93.4, 97.6)
Total	15.5	(12.9, 18.6)	22.1	(18.6, 25.9)	3.8	(2.7, 5.4)	74.1	(70.9, 86.0)
Rural								
Male	25.6	(12.3, 30.4)	36.1	(31.1, 41.4)	4.9	(3.5, 6.6)	59.0	(55.5, 65.9)
Female	1.9	(0.8, 4.8)	2.2	(1.0, 5.1)	0.3	(0.1, 0.9)	97.5	(92.3, 98.3)
Total	13.8	(11.3, 16.8)	19.2	(16.3, 22.5)	2.1	(1.6, 2.8)	78.7	(75.8, 82.0)
Combined								
Male	26.5	(23.2, 30.1)	37.1	(33.3, 41.1)	6.0	(4.7, 7.6)	59.1	(55.1, 63.0)
Female	1.9	(0.9, 3.7)	2.6	(1.5, 4.4)	0.3	(0.2, 0.7)	97.1	(95.1, 98.3)
Total	14.3	(12.3, 16.5)	20.0	(17.7, 22.5)	2.6	(2.1, 3.2)	77.4	(75.4, 80.3)

Table 3.2.2 Percentage of drinkers (past 12 months) according to the frequency of drinking, mean number of standard drinks per day and pattern of drinking in the last seven days by gender and place of residence, Andhra Pradesh, 2007- 08

Alcohol consumption	Residence						Combined		
	Urban			Rural					
	Male	Female	Total	Male	Female	Total	Male	Female	Total
Frequency of drinking in past 12 months									
5-7 days per week	11.6	9.2	11.4	20.0	40.2	21.1	17.4	28.4	18.1
1-4 days per week	20.8	4.6	19.6	23.3	33.8	23.9	22.5	22.6	22.5
1-3 days per month	37.0	24.1	36.0	29.3	3.8	27.8	31.7	11.5	30.4
Less than once per month	30.6	62.1	33.1	27.4	22.2	27.1	28.4	37.4	29.0

Mean number of drinks on a drinking day	2.8	2.3	2.7	3.4	1.4	3.3	3.2	1.7	3.1
Drinks during last 7 days									
Alcohol consumed on 4+days (%)	15.7	21.1	16.0	32.7	62.5	34.8	27.4	51.7	29.0
**Binge drinking on any day (%)	18.6	31.2	19.3	26.5	3.1	24.9	24.0	10.5	23.2
20+ drinks in 7 days (%)	8.0	21.1	8.7	19.2	4.6	18.2	15.7	8.9	15.2
Average standard drinks per day	1.2	1.7	1.2	1.9	0.9	1.8	1.7	1.1	1.6

** 5+ drinks on any day for male; and 4+drinks on any day for female

percent of such respondents in urban area and 24% respondents in rural area consumed alcohol 1-4 days every week in past one year. And 17% of the males and 28 % of females were consumed alcohol 5-7 days per week in the past one year. The average numbers of drinks consumed on a drinking day was 3 drinks.

The respondents who were current drinker were also asked about their behaviour in terms of the number of days and number of drinks per day they took in the past 7 days preceding the survey. The survey found that 16% respondents of urban, 35% respondents of rural and 29% in combined sample respondents consumed alcohol at least 4 days a week. About 15% of the respondents had 20 or more drinks in the last 7 days. About 19% of current drinkers of urban and 25% of rural were in high risk drinking (binge drinking). The average standard drink consumed per day is calculated using the data collected

for alcohol consumption by current drinkers in the week preceding the survey which was about two drinks.

Table 3.2.3 presents the mean age of initiation of alcohol use by sex and the place of residence for current and past drinkers in the age group 15-34 years and 35-64 years. The mean age of initiation of alcohol consumption regularly in the age group of 15-34 years was 20 years irrespective of the sex and place of residence. The mean age of initiation in the age group 35-64 years was 25 years.

Table 3.2.4 presents the percentage of current daily smokers and smokeless tobacco users and current drinkers by age, education and occupation. The percentage of daily smoker was high in the age group 35-44 years (28%), among illiterates (58%) and agriculturist (42%). Similarly, percentage of smokeless tobacco users was high in the age group

Table 3.2.3 Mean age of initiation of alcohol use by sex and place of residence, Andhra Pradesh, 2007-08

Residence/ Sex	Alcohol users			
	Age of Initiation (15-34 years)		Age of Initiation (35-64 years)	
	Mean	95% CI	Mean	95% CI
Urban				
Male	20	(19.5, 21.5)	25	(24.5, 25.5)
Female	*	*	20	(19.5, 28.5)
Total	20	(18.5, 20.5)	25	(24.5, 25.5)
Rural				
Male	20	(19.5, 20.5)	23	(21.5, 25.5)
Female	*	*	18	(15.5, 20.5)
Total	20	(18.5, 20.5)	21	(20.5, 23.5)
Combined				
Male	20	(19.5, 20.5)	25	(24.5, 25.5)
Female	*	*	20	(18.5, 20.5)
Total	20	(19.5, 20.5)	25	(23.5, 25.5)

* Figure not shown; based on fewer than 15 unweighted cases.

Table 3.2.4 Percentage of current daily smokers, daily smokeless tobacco users and current drinkers across age, education and occupation, Andhra Pradesh, 2007- 08

Characteristic	Smoker	Smokeless tobacco user	Current drinkers
Age group			
15-24	9.6	14.6	12.9
25-34	21.5	29.1	28.6
35-44	27.8	24.6	29.3
45-54	24.9	17.2	19.5
55-64	16.2	14.5	9.7
Total	100.0	100.0	100.0
Education			
Illiterate	57.7	58.2	50.2
Primary	12.2	7.6	10.0
Middle	12.0	11.4	14.2
Secondary	11.5	13.2	13.9
Higher Secondary	3.0	7.1	5.5
College & above	3.6	2.5	6.2
Total	100.0	100.0	100.0
Occupation			
Executive/Business	9.1	8.7	11.9
Agriculture	42.3	37.2	39.3
Domestic Work	4.3	9.3	3.2
Services/Sales	4.6	5.3	6.7
Manual Worker	32.8	32.6	33.3
Other	6.9	6.9	5.6
Total	100.0	100.0	100.0
Number (n)	1105	619	1220

25-34 years (29%), among illiterates (58%) and agriculturists (37%). The percentage of current drinkers was high (29%) in the age group 35-44, among illiterates (50%) and those who reported their occupation as agriculture (39%).

Interesting observations of alcohol consumption was that about 37% of men consumed alcohol at least once in last one year whereas 27% of men in last one month. The alcohol consumption among females was very low. Those who consumed alcohol in last seven days, 23% of them were binge drinkers. The mean age of initiation of alcohol consumption by young age (15-34 years) men was 20 years.

3.3 FRUITS AND VEGETABLES CONSUMPTION

Survey asked questions about the number of days in a typical week on which fruits and vegetables are consumed by the respondents and the number of servings of fruits and vegetables consumed on one of those days.

Table 3.3.1 presents mean numbers of days in a week fruits and vegetables consumptions by sex and place of residence.

Mean number of days in a week people in Andhra Pradesh consumed vegetables and fruits was 5 days and 2 days a week respectively. The mean

Table 3.3.1 Mean number of days in a week fruit and vegetable consumed by the respondents according to gender and place of residence, Andhra Pradesh, 2007- 08

Fruits and vegetables consumption per week	Residence						Combined		
	Urban			Rural					
	Male	Female	Total	Male	Female	Total	Male	Female	Total
Mean number of days fruits consumed	2.7	2.4	2.5	1.8	1.4	1.6	2.0	1.7	1.9
Mean number of days vegetables consumed	5.0	5.0	5.0	4.9	4.9	4.9	5.0	4.9	4.9
Less than five servings of fruits & vegetables consumed per day (%)	82.2	85.8	84.0	87.5	91.8	89.6	86.0	90.1	88.0

number of days when fruits consumed was high in case of urban (3 days) as compared to that in rural (2 days) population.

Eighty eight percent of respondents report that they had less than five servings of fruits and vegetables per day on those days when they consumed it. It was 84% among urban and 90% among rural respondents. From Table 3.3.2, it can be seen that the mean number of servings of fruits and vegetables in one particular day was 2 servings across the sex and place of residence.

Nutritional inadequacy is the major risk factor of many non-communicable diseases. Overall, 88% of population in Andhra Pradesh consumed less than five servings of fruits and vegetables per day, which was inadequate as per WHO recommended standards. On an average only two days in a week people consumed fruits against vegetables consumed on 5 days. This is

an important health issues and needs to be address with more emphatically.

Food and Oil Consumption

The percentage of respondents according to the intake of specific food items at least once a week by sex and place of residence is provided in Table 3.3.3. The specific food items included cheese/butter, fried local foods, red meat, eggs, chicken, aerated soda, sweetened drinks, pizza/burger/French fries, cakes/pastries or other bakery items, chips/*namkeen* etc. in the survey. About 70% population consumed eggs, 50% consumed chicken, 28% consumed fish, 34% consumed red meat, 15% consumed fried local foods and 3% consumed cakes/pastries or other bakery items at least once a week. The daily consumption of these items were very less, not more than 4% of the population.

Table 3.3.2 Mean number of servings of fruits, vegetables consumed in one particular day by gender and place of residence Andhra Pradesh, 2007- 08

Mean Number of servings of fruits/vegetables/both in a day	Residence						Combined		
	Urban			Rural					
	Male	Female	Total	Male	Female	Total	Male	Female	Total
Servings of fruit	0.8	0.7	0.8	0.6	0.5	0.6	0.7	0.6	0.6
Servings of vegetable	1.5	1.4	1.4	1.3	1.2	1.3	1.4	1.3	1.3
Servings of fruit and vegetable	2.4	2.1	2.3	2.0	1.8	1.9	2.1	1.9	2.0

Table 3.3.3 Percentage of respondents according to the intake of specific food items by place of residence, Andhra Pradesh, 2007- 08

Specific Food Items	Residence				Combined	
	Urban		Rural			
	Daily	At least once in a week	Daily	At least once in a week	Daily	At least once in a week
Cheese/ Butter	4.9	7.0	2.1	5.3	2.9	5.8
Fried local foods	4.7	20.2	2.0	12.5	2.8	14.7
Red Meat	0.9	41.0	0.1	31.2	0.3	34.0
Eggs	3.8	69.8	1.3	70.2	2.0	70.1
Chicken	0.3	58.0	0.5	47.3	0.5	50.4
Fish	0.2	22.9	2.4	30.2	1.8	28.1
Aerated Soda	0.8	9.6	0.7	7.5	0.7	8.1
Sweetened drinks	1.7	11.6	0.2	3.5	0.6	5.8
Pizza/ burgers/ French fries etc.	0.2	3.2	0.7	0.9	0.0	1.4
Cakes, Pastries or other bakery items	0.9	7.2	0.3	1.7	0.4	3.3
Chips, <i>Namkeen</i> etc.	2.4	24.2	2.2	15.4	2.3	23.4

Table 3.3.4 presents the type of edible oil used for cooking by the sample households in rural and urban residence. It shows that the use of palm oil for cooking was highest (49% households with 30% in urban and 56% in rural households) followed by groundnut oil (29% households with 35% in urban and 27% in rural households), sunflower oil (19% household with 33% in urban and 13% in rural households).

Table 3.3.4 Percentage of households according to type of oil consumption, Andhra Pradesh, 2007-08

Type of oil	Residence		Combined
	Urban	Rural	
Coconut oil	0.1	0.0	0.0
Groundnut oil	34.6	27.3	29.4
Sunflower oil	32.6	12.9	18.5
Soyabean oil	0.1	0.4	0.3
Palm oil	29.8	56.1	48.6
Vanaspati oil	0.2	0.3	0.3
Others	2.6	3.0	2.9
Total	100.0	100.0	100.0

3.4 PHYSICAL ACTIVITY

It is well known that lack of physical activity leads

to obesity, hyperlipidemia, diabetes mellitus, hypertension and coronary heart disease. An account of physical activities of respondents in terms mean time spent (in minutes) in doing physical activity at work, while traveling for work and recreation by sex and the place of residence, is provided in Table 3.4.1. On an average, people in Andhra Pradesh were doing some physical activity for a duration of 1210 MET minutes per day, 1034 MET minutes per day in urban and 1280 MET minutes per day in rural area. Men, on an average, spent 1394 MET minutes a day while women spent 1022 MET minutes a day on physical activity. The mean time spent in work related physical activity by the respondents was 279 minutes per day which was 206 minutes per day for urban and 238 minutes per day for rural respondents. The time spent in work related physical activity was more among men (239 minutes/day) than women (218 minutes/day).

The mean time spent in travel related activity (cycling/walking) was found to be 31 minutes per day (26 minutes in urban and 33 minutes in rural areas). It was more among men (38 minutes per day) as compared to women (24 minutes per day). The survey also reports that the mean time spent in recreational activities which was low (4 minutes per day) 6minutes for urban and 3 minutes for rural respondents). Men spent more time

Table 3.4.1 Mean time spent (in minutes) on physical activity per day by sex and residence, Andhra Pradesh, 2007- 08

Physical Activity	Residence						Combined		
	Urban			Rural					
	Male	Female	Total	Male	Female	Total	Male	Female	Total
Total Physical Activity (MET)									
Mean	1202.2	857.5	1034.0	1471.9	1085.7	1279.5	1384.0	1021.7	1209.6
95% CI Lower	1109.0	814.0	973.6	1361.6	1001.7	1188.3	1312.4	962.6	1143.3
Upper	1295.4	901.0	1094.3	1582.2	1169.8	1370.7	1475.5	1080.7	1275.8
Work Related Activity									
Mean	222.2	188.0	205.5	246.0	229.6	237.8	239.1	217.9	228.6
95% CI Lower	203.7	178.7	193.3	227.8	213.3	221.8	225.4	206.4	216.8
Upper	240.8	197.3	217.7	264.1	245.9	253.9	252.8	229.5	240.4
Travel Related Activity									
Mean	33.3	18.8	26.3	39.4	26.4	32.9	37.7	24.3	31.0
95% CI Lower	30.2	16.5	23.9	34.7	22.2	34.2	34.2	21.3	28.0
Upper	36.4	21.2	28.6	44.2	30.5	41.1	41.1	27.2	34.1
Recreational Activity									
Mean	9.3	1.8	5.6	5.6	3.0	6.7	6.7	0.7	3.7
95% CI Lower	7.3	1.1	4.5	3.2	1.8	4.9	4.9	0.5	2.8
Upper	11.3	2.4	6.8	8.1	4.2	8.5	8.5	1.0	4.6
Number	1257	1493	2750	1462	3468	2719	2719	3499	6218

(7 minutes per day) than women (less than one minutes per day) in recreational activities.

According to WHO Global Physical Activity Questionnaire Analysis Guidelines⁷, the total physical activity of the respondents is classified under three categories low, medium and high on the basis of duration for which they perform physical activities of varying intensity. The percentage of respondents according to three categories of physical activity by sex and place residence is presented in Table 3.4.2. Majority of the respondents (68% overall, 78% in urban and 64% in rural) reported low physical activity, 28% of respondents (21% in urban and 31% in rural) reported medium physical activity and only 4% of

respondents (2% in urban and 5% in rural) reported a high level of physical activity.

Table 3.4.3 presents the percentage of respondents according to their category of time spent in physical activity by age and sex. About 2% of old age (55-64) respondents was recorded in high level of physical activity which was less than 6% in the younger age groups of people in Andhra Pradesh. The total time spent daily in sedentary activities is also recorded and provided in Table 3.4.4. Majority of the respondents (33%) spent 2-3 hours in sedentary activities followed by 23% of respondents spent more than 4 hours and 21% of the respondents spent 1-2 hours.

Table 3.4.2 Percentage of respondents classified in three categories of total physical activity per day (P & 95% CI), by sex and place of residence, Andhra Pradesh, 2007-08

Physical Activity	Residence						Combined		
	Urban			Rural					
	Male	Female	Total	Male	Female	Total	Male	Female	Total
Low	64.3	91.3	77.5	52.5	75.2	63.8	55.9	79.7	67.7
95% CI Lower	59.4	88.9	74.4	46.5	70.4	59.0	51.4	76.3	64.2
95% CI Upper	69.0	93.2	80.3	58.4	79.4	68.3	60.3	82.7	71.0
Medium	32.7	8.3	20.8	38.3	23.4	30.9	36.7	19.2	28.0
95% CI Lower	28.5	6.4	18.3	33.4	19.5	27.1	33.0	16.3	25.2
95% CI Upper	37.2	10.6	23.6	43.4	28.0	35.0	40.5	22.4	31.0
High	2.9	0.4	1.7	9.2	1.4	5.3	7.4	1.1	4.3
95% CI Lower	1.6	0.2	0.9	6.9	0.7	3.9	5.7	0.6	3.3
95% CI Upper	5.3	1.1	3.1	12.2	2.9	7.2	9.6	2.2	5.6

Table 3.4.3 Percentage of respondents (with 95% confidence interval) according to three categories of total physical activity by age group and sex, Andhra Pradesh, 2007-08

Age group	Sex								
	Men			Women			Both Sex		
	Low	Medium	High	Low	Medium	High	Low	Medium	High
15-24	62.0 (56.0,67.6)	30.8 (25.7,36.3)	7.3 (4.7,11.1)	84.0 (79.1,87.9)	15.5 (11.7,20.3)	0.5 (0.2,1.7)	72.7 (68.3,76.7)	23.2 (19.7,27.4)	4.0 (2.6,6.0)
25-34	52.5 (46.3,58.6)	41.1 (35.5,46.9)	6.5 (4.1,10.0)	73.8 (68.1,78.7)	24.4 (19.4,29.3)	2.2 (1.1,4.3)	63.3 (58.9,67.5)	32.4 (28.5,36.5)	4.3 (2.9,6.4)
35-44	48.1 (41.8,54.4)	41.6 (36.4,47.0)	10.3 (7.5,14.1)	76.0 (70.2,81.0)	22.5 (17.6,28.2)	1.5 (0.5,4.2)	61.6 (56.2,66.8)	32.3 (27.9,37.1)	6.1 (4.2,8.6)
45-54	50.6 (44.7,56.5)	42.6 (37.1,48.2)	6.8 (4.0,11.3)	77.2 (71.4,82.1)	22.7 (17.8,28.5)	0.1 (0.0,0.9)	63.4 (59.3,67.3)	33.0 (29.7,36.5)	3.6 (2.1,6.1)
55-64	72.6 (66.0,78.4)	23.0 (17.6,29.3)	4.4 (2.6,7.3)	93.8 (91.5,95.5)	5.7 (4.0,7.9)	0.5 (0.2,1.7)	83.8 (80.4,86.8)	13.8 (11.1,17.1)	2.3 (1.4,3.8)

Note: WHO Steps guidelines used to calculate the cut off value of low, medium and high for total physical activity.

Table 3.4.4 Sex wise percentage of respondents classified according to total time spent in sedentary activity per day by type of residence, Andhra Pradesh, 2007-08

Time spent sitting/ reclining	Residence						Combined		
	Urban			Rural					
	Male	Female	Total	Male	Female	Total	Male	Female	Total
Less than 1 hours	2.7	1.2	2.0	5.8	3.8	4.8	4.9	3.1	4.0
1-2 hours	15.0	8.2	11.7	26.5	23.8	25.1	23.1	19.4	21.3
2-3 hours	32.7	23.0	28.0	37.1	31.6	34.3	35.8	29.1	32.5
3-4 hours	19.3	23.7	21.5	15.6	19.7	17.6	16.6	20.8	18.7
More than 4 hours	30.2	23.7	36.9	15.0	21.2	18.1	19.4	27.5	23.4

3.5 SOCIO-DEMOGRAPHIC PATTERN

The socio-demographic pattern of non-communicable disease risk factors (tobacco, alcohol, fruits and vegetable consumption, and physical activity) are presented in Table 3.5.1 for urban, Table 3.5.2 for rural and Table 3.5.3 for combined (all respondents).

Tobacco

Tobacco is mainly used either in the form of smoking or non-smoking (chewing with lime or *Pan*) among urban and rural residents. The prevalence and pattern of smoking among urban male respondents was increasing with age (9% in 15-24 to 40% in 45-54). But, it was decreasing with increasing level of education (45% among illiterates to 10% in higher secondary). Prevalence of smoking among female respondents was very low compare with males across all the categories. Occupation is an important socioeconomic indicator and differences in pattern of smoking tobacco was observed from one category of occupation to another. Prevalence of smoking among the people working in agriculture (43%) and manual worker (33%) was high compare with other categories of occupation. A similar pattern of increasing in prevalence with age and decreasing with level of education was also observed with smokeless tobacco users. The prevalence of smokeless tobacco users among urban respondents was also increasing with age (6% in 15-24 to 11% in 55-64 aged respondents). Prevalence of smokeless tobacco users among females was recorded low (3%), but the pattern was increasing with age. However, the prevalence with education was showing a declining pattern with increasing level of education (12% among illiterates to 6% in higher Secondary). Prevalence of smokeless tobacco users among agriculture (14%) and manual worker (21%) was high compare with other categories of occupation. Among urban male respondents, smokers were higher (24%) than smokeless tobacco users (15%), but the pattern was similar in both.

Rural-urban differences in the prevalence of smoking and smokeless tobacco users were observed across all the socio-demographic categories. Prevalence of smoking among rural male respondents was high (35%) compare with urban males (24%). Overall, pattern of smoking and smokeless tobacco use in urban and rural subgroups of population remain similar across age, education and occupation. The prevalence of smoking among rural male respondents shows the increasing pattern with age (12% in 15-24 to 57% in 55-64 aged). Prevalence of smoking among agriculture (42%) and manual worker (39%) of male rural respondents was high compare with other categories of occupation. Similar pattern of smoking and smokeless tobacco users was observed in the combined population (Table 3.5.3).

Alcohol

The prevalence of current alcohol drinkers among urban male respondents was high among the adults (49% in 25-34, 52% in 35-44, 42% in 45-54 and 32% in 55-64 age groups). It was comparatively low among younger age (25% in 15-24). The increasing pattern of alcohol prevalence with age was observed up to age group of 35-44 and than the pattern declined. Prevalence of drinking alcohol among male respondents was recorded high in lower level of education i.e. illiterates (52%) and Primary (56%) and than started decline among higher (30% among collage) level. Prevalence of drinking alcohol among manual worker (55%), agriculture (52%) and Service (42%) categories of occupation of urban male respondents was high. Low prevalence of drinking alcohol was recorded (3.5%) among the female urban respondents (Table 3.5.1). Among the rural male respondents, current alcohol users were high in the adult age groups (37% in 25-34, 50% in 35-44, 48% in 45-54 and 39% in 55-64 age groups). A similar pattern of prevalence of alcohol use was observed in the rural population across age, education and occupation (Table 3.5.2).

Table 3.5.1 Percentage of respondents in the category of some high risk factors of NCD (current daily smokers, daily smokeless tobacco user, current drinkers, low fruits and vegetables intake and low physical activity) across age, education, occupation and sex, urban, Andhra Pradesh, 2007- 08

Characteristic	Smoker			Smokeless tobacco user			Current drinkers			Less than five servings of fruits & vegetables consumed per day			Low physical activity		
	Male	Female	Total	Male	Female	Total	Male	Female	Total	Male	Female	Total	Male	Female	Total
Age group															
15-24	9.1	0.0	4.6	9.7	1.1	5.5	24.6	3.0	14.0	82.2	84.8	83.5	75.7	93.4	84.4
25-34	26.2	0.0	13.1	20.0	1.5	10.7	49.3	2.6	26.0	77.7	82.9	80.3	56.6	88.5	72.5
35-44	29.6	0.3	15.6	19.3	4.4	12.2	52.4	4.3	29.4	83.4	86.9	85.1	55.7	88.4	71.3
45-54	40.1	3.4	23.2	13.5	5.6	9.9	42.2	5.4	25.3	86.2	88.8	87.4	54.9	92.0	71.9
55-64	34.3	4.1	18.9	13.7	8.9	11.3	31.7	3.8	17.4	87.4	92.9	90.2	84.3	98.1	91.4
Total	23.9	0.8	12.6	15.2	3.0	9.2	39.7	3.5	22.1	82.2	85.8	84.0	64.3	91.3	77.5
Education															
Illiterate	45.1	2.6	14.9	23.1	6.9	11.6	51.7	7.4	20.3	83.1	90.1	88.1	45.1	84.3	72.9
Primary	48.3	0.0	22.1	24.8	5.2	14.1	56.1	3.6	27.6	89.5	83.3	86.1	48.9	94.7	73.8
Middle	35.0	0.0	18.1	20.4	0.8	10.9	49.1	3.1	26.9	80.4	86.7	83.4	56.1	92.5	73.6
Secondary	16.0	0.0	8.7	17.5	0.3	9.6	36.8	0.8	20.3	83.9	80.1	82.1	63.8	93.8	77.6
Higher Secondary	10.2	0.0	6.3	9.8	0.0	6.1	31.0	1.7	19.9	81.8	85.9	83.3	76.6	96.2	84.0
College & above	12.5	0.0	8.6	5.6	0.0	3.9	29.6	0.0	20.4	78.6	85.8	80.9	77.5	95.8	83.2
Total	23.9	0.8	12.6	15.2	3.0	9.2	39.7	3.5	22.1	82.2	85.8	84.0	64.3	91.3	77.5
Occupation															
Executive/Business	19.9	0.0	16.4	13.8	2.9	11.9	36.8	4.2	31.1	82.4	84.9	82.8	65.9	75.2	67.5
Agriculture	42.7	3.1	25.3	22.9	1.4	13.5	52.0	4.0	30.9	86.1	83.1	84.8	50.6	58.8	54.2
Domestic Work	**	0.6	0.6	**	2.7	2.7	**	2.9	2.9	**	85.9	86.0	**	97.8	97.9
Services/Sales	25.3	0.9	19.5	9.8	3.9	8.4	41.9	5.6	33.3	79.6	84.4	80.8	62.5	75.6	65.7
Manual Worker	33.2	2.4	27.3	24.6	7.7	21.4	54.9	9.5	46.3	79.6	90.1	81.6	44.5	65.2	48.5
Other	11.2	0.7	8.0	4.5	1.7	3.7	18.2	1.8	13.2	86.0	84.5	85.6	94.7	97.5	95.5
Total	23.9	0.8	12.6	15.2	3.0	9.2	39.7	3.5	22.1	82.2	85.8	84.0	64.3	91.3	77.5
Number (n)	1257	1493	2750	1257	1493	2750	1257	1493	2750	1257	1493	2750	1257	1493	2750

** Figure not shown; based on fewer than 15 unweighted cases

Fruits and Vegetables

Though fruits and vegetable consumption reduces the risk of non-communicable diseases, but the survey showed larger proportion of population consumed inadequate amount of fruits and vegetables (i.e. less than five servings of fruits and vegetables per day). Prevalence of low (inadequate) consumption was recorded high (84%) among urban population which varies with age groups (80% in 25-34 to 90% in 55-64). Inadequate consumption of fruits and vegetables by education was high among illiterate (88%) and it was found declining with increasing level of education (81% among College). Differences in occupation categories were also observed. It was varying between 81% among Service class to 86% among domestic worker in urban population (Table 3.5.1). A similar pattern of inadequate consumption of fruits and vegetables was observed among rural population. Prevalence of inadequate consumption was varying with age groups (87% in 25-34 to 95% in 55-64). A declining pattern with increasing level of education was reflected (94% among illiterates, 83% among higher education level). In the occupational categories, inadequate consumption of fruits and vegetables was high among the people working as domestic (92%) and agriculture (90%). It was comparatively low (78%) among executive category (Table 3.5.2). Overall, prevalence and pattern of consumption of fruits and vegetables by age, education and occupation was similar as recorded in rural and urban population of Andhra Pradesh (Table 3.5.3).

Physical Activity

The differences in the prevalence of low physical activity were recorded across age, sex, education and occupation in urban population (Table 3.5.1). Large proportion of urban respondents were recorded in the

category of low physical activity (78%) and it was varying with age groups (84% in 15-24, 73% in 25-34, 71% in 35-44, 72% in 45-54 and 91% in 55-64). Prevalence of low physical activity was high (91%) among old age people (55-64) as compare with younger age (84%). Gender difference of low physical activity was recorded high (91%) among female respondents compare with males (64%), and such differences remain across all the age groups (Table 3.5.1). The pattern of low physical activity by educational categories was increasing with level of education especially among male respondents (45% of illiterates to 77% of higher level). Accordingly, low physical activity was recorded high among the domestic (98%), executive and business (68%) categories of occupation. Those working in agriculture and manual worker were doing more physical activity (Table 3.5.1).

Urban-rural comparison of low physical activity demonstrated that rural population (64%) was doing more physical work than urban (78%) and such differences observed across all age groups and sex (Table 3.5.2). Low physical activity by education was observed more among higher level of education (85%) compare with lower level (60% among illiterates) in rural population. Similarly, occupational differences in low physical activity were also observed across all the categories. The occupation of people working as agriculture and manual worker were doing more physical work compare with others. Low physical activity was high among the domestic worker (97%) in the rural and urban population (Table 3.5.3).

Physical inactivity is one of the important risk factors of NCD. Most important point to be noted that seven out of ten individual adult population was categorized into low level of physical activity. This invites special attention to health planner.



Table 3.5.2 Percentage of respondents in the category of some high risk factors of NCD (current daily smokers, daily smokeless tobacco user, current drinkers, low fruits and vegetables intake and low physical activity) across age, education, occupation and sex, rural, Andhra Pradesh, 2007- 08

Characteristic	Smoker			Smokeless tobacco user			Current drinkers			Less than five servings			Low physical activity of fruits & vegetables consumed per day		
	Male	Female	Total	Male	Female	Total	Male	Female	Total	Male	Female	Total	Male	Female	Total
Age group															
15-24	11.9	0.8	6.5	8.2	0.0	4.2	18.7	0.5	9.9	85.7	90.8	88.2	56.0	79.8	67.5
25-34	29.9	2.1	15.6	18.0	2.2	9.9	37.1	1.1	18.7	84.5	88.9	86.8	50.8	67.9	59.6
35-44	49.1	4.9	27.5	13.9	7.2	10.6	50.1	3.2	27.2	85.7	92.3	88.9	45.0	71.3	57.9
45-54	53.5	12.7	33.6	11.7	10.7	11.2	48.0	5.4	27.2	94.1	95.5	94.8	49.0	72.1	60.3
55-64	57.1	15.6	35.0	15.3	14.8	15.0	38.6	3.4	19.8	94.6	95.7	95.2	68.9	92.5	81.5
Total	34.5	5.2	19.9	13.0	5.1	9.1	36.1	2.2	19.2	87.5	91.8	89.6	52.5	75.2	63.8
Education															
Illiterate	52.1	8.9	24.7	16.2	8.7	11.4	49.4	3.6	20.3	91.8	94.7	93.6	44.5	69.5	60.2
Primary	42.7	0.4	23.3	14.8	0.9	8.4	43.8	0.2	23.8	87.3	92.0	89.4	41.8	76.6	57.8
Middle	28.0	0.0	15.5	11.6	0.0	6.4	30.5	0.0	16.9	86.3	86.7	86.5	44.9	78.5	59.9
Secondary	22.0	0.6	14.6	11.1	0.0	7.2	25.7	1.3	17.2	84.0	87.0	85.1	61.2	91.6	71.8
Higher Secondary	10.4	0.0	7.4	13.2	0.0	9.4	21.7	0.0	15.6	84.2	80.9	83.2	72.4	96.7	79.2
College & above	9.3	0.0	7.3	1.7	0.0	1.4	12.4	0.0	9.7	83.3	79.9	82.5	82.3	93.6	84.7
Total	34.5	5.2	19.9	13.0	5.1	9.1	36.1	2.2	19.2	87.5	91.8	89.6	52.5	75.2	63.8
Occupation															
Executive/Business	25.4	5.9	19.7	6.5	4.6	6.0	24.9	3.1	18.5	77.5	77.6	77.5	61.8	64.9	62.7
Agriculture	42.1	5.4	26.2	15.4	5.8	11.2	44.0	2.2	25.9	88.6	91.9	90.1	40.2	54.1	46.2
Domestic Work	**	4.5	4.6	**	3.6	3.6	**	2.2	2.2	**	92.3	92.4	**	96.6	96.7
Services/Sales	20.4	1.6	15.9	10.4	10.5	10.4	29.1	0.0	22.2	81.6	87.1	82.9	63.5	92.6	70.3
Manual Worker	39.3	6.9	29.2	15.2	6.9	12.6	38.8	3.3	27.7	86.5	93.7	88.7	46.0	46.5	46.1
Other	14.2	6.9	12.1	6.6	7.4	6.8	17.4	1.0	12.6	93.3	92.3	93.0	90.4	93.0	93.0
Total	34.5	5.2	19.9	13.0	5.1	9.1	36.1	2.2	19.2	87.5	91.8	89.6	52.5	75.2	63.8
Number (n)	1462	2006	3468	1462	2006	3468	1462	2006	3468	1462	2006	3468	1462	2006	3468

** Figure not shown; based on fewer than 15 unweighted cases

Table 3.5.3 Percentage of respondents in the category of some high risk factors of NCD (current daily smokers, daily smokeless tobacco user, current drinkers, low fruits and vegetables intake and low physical activity) across age, education, occupation and sex, combined, Andhra Pradesh, 2007-08

Characteristic	Smoker		Smokeless tobacco user		Current drinkers		Less than five servings		Low physical activity of fruits & vegetables consumed per day							
	Male	Female	Male	Female	Male	Female	Male	Female	Male	Female	Total					
Age group																
15-24	11.0	0.6	5.9	0.3	8.6	0.3	4.6	20.5	1.3	11.1	84.6	88.9	86.7	62.0	84.0	72.7
25-34	28.8	1.5	14.9	2.0	18.6	2.0	10.1	40.7	1.6	20.8	82.5	87.2	84.9	52.5	73.8	63.3
35-44	43.5	3.7	24.1	6.4	15.4	6.4	11.1	50.8	3.5	27.8	85.1	90.8	87.8	48.1	76.0	61.6
45-54	49.8	10.3	30.8	9.4	12.2	9.4	10.8	46.4	5.4	26.7	91.9	93.8	92.8	50.6	77.2	63.4
55-64	51.5	13.0	31.2	13.4	14.9	13.4	14.1	36.9	3.5	19.3	92.9	95.1	94.0	72.6	93.8	83.8
Total	31.5	4.0	17.9	4.5	13.6	4.5	9.1	37.1	2.6	20.0	86.0	90.1	88.0	55.9	79.7	67.7
Education																
Illiterate	51.2	7.8	23.1	17.1	8.4	8.4	11.4	49.7	4.3	20.3	90.7	93.9	92.8	44.5	71.9	62.2
Primary	43.8	0.3	23.0	16.7	2.0	2.0	9.7	46.2	1.0	24.6	87.7	89.8	88.7	43.2	81.2	61.3
Middle	30.1	0.0	16.3	14.2	0.3	0.3	7.8	36.1	1.0	20.0	84.6	86.7	85.5	48.2	83.1	64.2
Secondary	20.1	0.3	12.5	13.1	0.1	0.1	8.1	29.2	1.1	18.3	84.0	84.1	84.0	62.0	92.5	73.8
Higher Secondary	10.3	0.0	6.9	11.7	0.0	0.0	7.9	25.8	0.9	17.6	83.1	83.6	83.3	74.2	96.4	81.5
College & above	11.2	0.0	8.1	4.0	4.0	0.0	2.9	22.2	0.0	16.1	86.6	83.9	81.5	79.6	95.1	83.8
Total	31.5	4.0	17.9	13.6	13.6	4.5	9.1	37.1	2.6	20.0	86.0	90.1	88.0	55.9	79.7	67.7
Occupation																
Executive/Business	22.1	3.3	17.9	10.9	10.9	3.9	9.3	32.0	3.6	25.5	80.4	80.8	80.5	64.3	69.3	65.4
Agriculture	42.1	5.3	26.2	15.7	15.7	5.6	11.3	44.0	2.3	26.1	88.5	91.5	89.8	40.7	54.3	46.6
Domestic Work	**	3.0	3.0	**	**	3.3	3.3	**	2.4	2.4	**	89.9	89.9	**	97.1	97.1
Services/Sales	22.6	1.3	17.5	10.1	10.1	7.5	9.5	34.9	2.5	27.3	80.7	85.8	81.9	63.1	84.8	68.2
Manual Worker	37.3	6.0	28.7	18.2	18.2	7.0	15.1	43.9	4.5	33.0	84.3	93.0	86.7	45.5	50.1	46.8
Other	13.1	4.5	10.0	5.8	5.8	5.2	5.6	17.7	1.3	12.8	90.6	89.3	90.2	92.0	98.6	93.9
Total	31.5	4.0	17.9	13.6	13.6	4.5	9.1	37.1	2.6	20.0	86.0	90.1	88.0	55.9	79.7	67.7
Number (n)	2719	3499	6218	2719	2719	3499	6218	2719	3499	6218	2719	3499	6218	2719	3499	6218

** Figure not shown; based on fewer than 15 unweighted cases

CHAPTER 4

Hypertension and Diabetes

This chapter focuses on the prevalence of hypertension and diabetes in the study population along with the information regarding history of hypertension and diabetes and the nature of treatment advised by the treating physician.

4.1 HYPERTENSION

The blood pressure is an important determinant of the risk of cardiovascular diseases, ischemic heart disease, congestive cardiac failure and renal failure. In the survey the blood pressure of the respondents was measured using automated blood pressure measuring instrument (OMRON®). Table 4.1.1 provides percentage of respondents with history of raised blood pressure, treatment and life style modification advised, seeking consultation and treatment from AYUSH by sex and place of residence. Over all 8% respondents, 7% men, 8% women were found to have been diagnosed hypertension by the health professional. In the urban area, the prevalence of hypertension was 10% with 10% among men and 11% among women. In rural area, 6% of males

and 7% of females were hypertensive.

Of those who were diagnosed hypertension, majority of them (67%) were taking the prescribed medicine. A higher percentage of women (68%) than men (65%) were taking medicine. The phenomenon appears to be more common in rural area where 65% women against 61% men take medicine after they were diagnosed with hypertension. Seventy one percent of those who were diagnosed hypertension, were advised dietary modification including low salt intake; 36% were advised to lose weight and 7% were advised to increase physical activity. If the respondent was a smoker, 14% were advised to quit smoking. The percentage of those who received dietary advice including low salt intake was slightly higher for women (72%) than for men (70%), more for rural (72%) than urban (70%) respondents.

Less than one-tenth (8%) of those who were diagnosed hypertensive, have consulted AYUSH with 6% in urban and 10% in rural. By sex, 6% such men against 10% women were consulting AYUSH. Among these respondents who had consulted AYUSH, three-quarter

Table 4.1.1 Percentage of respondents with history of raised blood pressure, treatment and lifestyle modification advised, seeking consultation and treatment from an AYUSH practitioner by gender and place of residence, Andhra Pradesh, 2007- 08.

Hypertension	Residence						Combined		
	Urban			Rural					
	Male	Female	Total	Male	Female	Total	Male	Female	Total
Hypertension diagnosed by health professional (all respondents)	9.7	10.5	10.1	5.5	7.4	6.4	6.7	8.3	7.5
Diagnosed Hypertensive									
Currently taking drugs	71.1	73.4	72.3	60.6	65.1	63.2	65.0	68.1	67.0
Advised dietary modification	70.0	70.0	70.0	69.4	73.8	72.0	69.6	72.4	71.2
Advised to lose weight	48.5	37.1	42.7	31.8	32.3	32.1	38.8	34.0	36.2
Advised to quit smoking	23.0	1.2	11.9	27.7	7.5	16.1	25.7	5.3	14.5
Advised to increase physical activity	53.6	40.9	47.2	26.0	33.8	30.5	37.6	36.4	37.4
Consulted AYUSH practitioner	6.7	5.9	6.3	5.4	12.7	9.6	5.9	10.3	8.3
Taking treatment from AYUSH practitioner	89.9	100	94.7	84.7	63.0	68.3	87.1	70.7	76.0

were taking the treatment from the AYUSH practitioner which was 95% in case of urban and 68% in case of rural.

Table 4.1.2 presents the mean systolic and diastolic blood pressure by sex and place of residence. In the survey population the mean systolic blood pressure was 123 mm Hg of mercury while mean diastolic blood pressure was 77 mm Hg of mercury. These averages were same for rural and urban respondents, both among males and females.

According to WHO STEPS guidelines, the population

is categorized into four categories namely, normal, pre-hypertensive, Stage-I hypertensive and Stage-II hypertensive on the basis of their blood pressure level⁸. In the present survey, this categorization is done after recording the resting blood pressure for each study subject. The upper and the lower limit of the systolic and diastolic blood pressure for each category has been given in Table 4.1.3.

Table 4.1.4 gives the percentage of respondents according to categories of hypertension by sex and place

Table 4.1.2 Mean Systolic and Diastolic blood pressure by sex and place of residence, Andhra Pradesh, 2007- 08

Blood Pressure	Residence						Combined		
	Urban			Rural			Male	Female	Total
	Male	Female	Total	Male	Female	Total			
Systolic blood pressure	126.5	120.0	123.2	125.0	120.0	122.5	125.4	120.0	123.0
95% CI Lower	125.0	119.0	122.3	124.0	119.0	121.4	124.4	119.0	121.0
Upper	127.6	121.0	124.1	126.3	121.3	123.6	126.4	121.0	123.5
Diastolic blood pressure	79.2	76.5	78.0	77.6	76.0	77.0	78.0	76.1	77.1
95% CI Lower	78.4	76.0	77.3	76.4	75.0	76.0	77.2	75.4	76.4
Upper	80.1	77.1	78.5	79.0	77.0	78.0	78.9	76.8	77.0

Table 4.1.3 Categories of Hypertension

Category	Systolic Blood Pressure (mm Hg)	Diastolic Blood Pressure (mm Hg)
Normal	<120 and	<80
Pre-hypertension	120-139 or	80-89
Stage-I hypertension	140-159 or	90-99
Stage-II hypertension	≥160 or	≥ 100

Table 4.1.4 Percentage of respondents according to category of hypertension by sex and place of residence (P & 95% CI), Andhra Pradesh, 2007- 08

Category of Hypertension	Residence						Combined		
	Urban			Rural			Male	Female	Total
	Male	Female	Total	Male	Female	Total			
Normal	26.9	49.2	37.7	33.3	49.1	41.1	31.5	49.1	40.2
95% CI Lower	23.1	45.5	34.5	29.0	45.3	37.5	28.2	46.2	37.4
Upper	31.1	52.8	41.1	37.9	52.9	44.7	35.0	52.0	42.9
Pre - hypertension	53.4	38.3	46.1	46.8	37.4	42.2	48.7	37.6	43.3
95% CI Lower	49.4	34.9	42.8	43.2	34.2	39.5	45.8	35.1	41.1
Upper	57.3	41.9	49.3	50.6	40.7	44.9	51.6	40.2	45.4
Stage-I hypertension	14.7	9.6	12.2	15.9	9.7	12.8	15.5	9.7	12.7
95% CI Lower	13.0	8.2	11.0	13.4	8.0	11.3	13.7	8.3	11.5
Upper	16.7	11.2	13.6	18.7	11.8	14.5	17.6	11.2	13.9
Stage-II hypertension	4.9	2.9	4.0	4.0	3.8	3.9	4.3	3.6	3.9
95% CI Lower	3.7	2.2	3.2	3.1	2.8	3.2	3.5	2.8	3.3
Upper	6.5	3.8	4.9	5.2	5.2	4.9	5.2	4.5	4.6

and residence. Over all, 40% respondents were normal, 43% were in the category of pre-hypertension, 13% in stage-I hypertension and only 4% were in stage-II hypertension. Among males, 32% were normal, 49% were in the category of pre-hypertension, 16% were in stage-I hypertension and only 4% were in stage-II hypertension. Among females 49% were normal, 38% pre-hypertension, 10% stage-I hypertension and 4% stage-II hypertension. The composition appears to be same for urban and rural respondents.

4.2 SOCIO-DEMOGRAPHIC PATTERN OF HYPERTENSION

Hypertension is a major NCD risk factors especially related to cardiovascular disease. The socio-demographic pattern of respondents in the category of hypertension (stage I & II) are presented in Table 4.2.

Among the urban population, the prevalence of hypertension was 16% and it was recorded as increasing with age (3% in 15-24, 13% in 25-34, 24% in 35-44, 37% in 45-54 and 40% in 55-64). The prevalence among male respondents was high (20%) compare with females (13%), but the increasing pattern with age was observed in both sexes (Table 4.2). The prevalence of hypertension by education was 22% among illiterate and 15% among higher level. However, the prevalence was high among executive and business (20%), manual worker (20%) and agriculture (19%) categories of occupation. Low prevalence of hypertension was recorded among the domestic workers(14%). Overall, prevalence among rural population was 17% and the pattern was increasing with age (6% in 15-24 to 39% in 55-64). Similarly, high prevalence was observed among illiterate (18%), which was showing a declining

Table 4.2 Percentage of respondents in the category of stage I & stage II hypertension across age, education, occupation and by sex and residence, Andhra Pradesh, 2007- 08

Characteristic	Stage I & II hypertensive								
	Urban			Rural			Combined		
	Male	Female	Total	Male	Female	Total	Male	Female	Total
Age group									
15-24	4.9	1.6	3.3	7.7	4.3	6.0	6.8	3.4	5.2
25-34	17.9	7.9	12.9	16.0	6.3	11.0	16.6	6.8	11.6
35-44	29.5	18.0	24.0	22.0	13.8	18.0	24.1	15.0	19.7
45-54	38.7	35.3	37.2	38.5	31.0	34.9	38.6	32.0	35.5
55-64	40.7	40.2	40.4	39.9	37.8	38.8	40.0	38.3	39.1
Total	19.7	12.5	16.2	19.9	13.5	16.7	19.8	13.2	16.6
Education									
Illiterate	26.8	19.6	21.8	20.5	16.9	18.2	21.3	17.4	18.8
Primary	23.7	13.0	18.1	20.7	12.1	16.8	21.3	12.3	17.1
Middle	21.7	13.1	17.6	20.4	10.8	16.1	20.8	11.5	16.5
Secondary	18.2	8.1	13.5	20.0	4.3	14.7	19.5	6.0	14.3
Higher Secondary	13.7	6.3	10.8	19.6	2.2	14.6	17.1	4.4	12.9
College & above	18.5	7.2	14.8	14.5	9.5	13.4	16.7	7.9	14.2
Total	19.7	12.5	16.2	19.9	13.5	16.7	19.8	13.2	16.6
Occupation									
Executive/Business	21.0	17.8	20.4	30.9	14.8	26.4	25.2	16.2	23.1
Agriculture	24.1	11.7	18.7	23.2	12.4	18.5	23.3	12.4	18.5
Domestic Work	**	13.2	13.1	**	14.4	14.7	**	13.9	14.1
Services/Sales	19.3	12.8	17.7	11.4	19.5	13.3	14.9	16.5	15.3
Manual Worker	22.2	9.8	19.9	15.5	11.9	14.4	17.6	11.5	15.9
Other	13.6	6.6	11.5	15.0	15.0	15.0	14.5	11.8	13.7
Total	19.7	12.5	16.2	19.9	13.5	16.7	19.8	13.2	16.6
Number (n)	1123	1335	2458	1257	1493	2750	2514	3213	5727

** Figure not shown; based on fewer than 15 unweighted cases

pattern with increasing level of education (13% in higher level). In the occupational category, the prevalence was high among executive and business class (26%), agriculture (19%) and it was low among services and sales (14%) category. Overall, prevalence of hypertension was 16% in Andhra Pradesh and pattern of prevalence was increasing with age.

Most striking observations of blood pressure measurements was that only around 40% of the adult population surveyed had normal blood pressure. While 43% of the adult population was categorized into pre-hypertension group, another 13% were found in stage I hypertension with the remaining 4% in stage-2. On the contrary, only 8% of population reported history of hypertension, which requires urgent attention for intervention.

4.3 DIABETES

Diabetes mellitus is an important marker of risk for the arterial disease of the coronary, cerebral and

peripheral arterial trees, and for micro vascular disease leading to blindness and renal failure. In the survey, the history pertaining to diabetes was elicited from the respondents. Table 4.3 deals with the percentage of respondents with history of raised blood sugar, a treatment and life style modification advises by sex and place of residence. Over all, 2% of respondents (4% in urban and 2% in rural) had reported having raised blood sugar level in past 12 months. This percentage was 3% in males and 2% in females. Among those who were diagnosed diabetes, 16% of the respondents (17% in urban and 16% in rural) were currently taking insulin; Eighty six percent of respondents (86% in urban and 85% in rural) were taking oral hypoglycemic drugs. A good proportion of respondents reported to have received advice from the treating physicians on their life style modification - 83% for dietary advice, 41% to reduce weight and 56% to increase physical activity.

Table 4.3. Percentage of respondents with history of raised blood sugar, treatment and lifestyle modification advised, seeking consultation and treatment from an AYUSH practitioner by gender and place of residence, Andhra Pradesh, 2007- 08

Blood sugar	Residence						Combined		
	Urban			Rural					
	Male	Female	Total	Male	Female	Total	Male	Female	Total
Raised blood sugar diagnosed (All respondents)	4.8	3.0	3.9	1.8	1.2	1.5	2.7	1.7	2.2
Diagnosed diabetics									
Currently taking insulin	16.5	17.3	16.8	17.1	13.2	15.6	16.8	15.2	16.2
Currently taking oral drugs	87.8	84.0	86.4	86.1	84.1	85.4	87.0	84.0	85.9
Advised dietary modifications	86.5	86.1	86.3	82.3	74.1	79.1	84.4	80.0	82.7
Advised to lose weight	49.5	50.0	49.7	33.5	30.6	32.4	41.7	40.2	41.1
Advised to increase physical activity	63.5	64.5	63.8	57.5	46.3	53.1	60.5	55.3	58.5
Consulted AYUSH practitioner	4.7	7.1	5.6	13.3	4.1	9.7	8.9	5.6	7.6
Taking treatment from AYUSH practitioner	100.0	56.0	79.0	59.2	100.0	66.0	70.3	72.3	70.9

CHAPTER 5

Physical Measurements

This chapter describes various physical measurements such as height, weight, waist circumference and body mass index (BMI), which are key indicators for surveillance of non-communicable diseases. Weight of an individual is directly related to the Body Mass Index (BMI), waist circumference, blood pressure and probability of developing diabetes mellitus-2.

5.1 WEIGHT

Having weight more than the ideal weight for age and height is a risk factor for development of colorectal cancer, uterine cancer, coronary artery disease and it would also exacerbate the symptoms of osteoarthritis. The weight is a continuous variable, reflecting the body mass of an individual in light clothing; it is used for calculating BMI.

5.2 HEIGHT

Height is another key variable required for calculation of body mass index (BMI). Height is a continuous variable measured with the individual standing on a firm leveled surface, without wearing any foot wear, and stand with feet together, with heels, calves, buttocks, dorsal spine and head in same plane.

5.3 BODY MASS INDEX (BMI)

BMI is a valid indicator for finding out whether the body weight of an individual is appropriate for the height of the individual. It is calculated from height and weight measurements as body weight per meter². Worldwide researches have shown that there is a strong association

between BMI and health risk. The excess of adipose tissue in the adults is associated with excess morbidity and mortality from a large number of health conditions like diabetes, hypertension, hypercholesterolemia, carcinomas of colon and breast, gall bladder stones and osteoarthritis. On the other hand low BMI is an indicator of risk to health, often being associated with tobacco, alcohol use and drug addiction (Table 5.1).

5.4 WAIST CIRCUMFERENCE

The waist circumference is one of the sensitive indicators for abdominal obesity. Abdominal obesity has got a stronger association with coronary heart diseases as compared to BMI. The waist measurement is taken at the level of mid point between the inferior margin of the rib and crest of ileum in the mid auxiliary plane, using a non-stretchable measuring tape, without clothing. A cut-off level of 102 cm. in males and 88 cm. in females have been recommended for developed countries (ATP3 Guidelines), however lower cut-off levels are appropriate for Indians- 90 cm in males and 80 cm in females (The Asia Pacific Guidelines)⁹.

Table 5.2 presents the BMI, mean height, weight and waist circumference by sex and the place of residence. The mean BMI in Andhra Pradesh was 21.7 kg/m² (23.1 for urban, 21.2 for rural, 21.8 for males and 21.7 for females). The mean height in the survey population was 158 centimeter (159 centimeter for urban, 157 centimeter for rural, 164 centimeter for males and 152 centimeter for females). The mean weight was 54 kg with 58 kg for urban, 52 kg for rural. By sex, the mean weight was 58 kg for males and 50 kg for females. The average waist circumference was 74

Table: 5.1 Categories of BMI

Body Mass Index (BMI)	Category of Relative Weight
<18.5	Under Weight
18.5- 24.9	Normal Weight
25.0- 29.9	Grade-1 Over Weight
30-39.9	Grade-2 Over Weight
≥ 40	Grade-3 Over Weight

Source: WHO Step-wise approach to NCD surveillance

Table 5.2 Mean value for body mass index (BMI), height, weight and waist circumference by gender and place of residence, Andhra Pradesh, 2007- 08

Physical Measurements	Residence						Combined		
	Urban			Rural					
	Male	Female	Total	Male	Female	Total	Male	Female	Total
BMI (kg/m ²)	22.9	23.2	23.1	21.3	21.1	21.2	21.8	21.7	21.7
95% CI Lower	22.3	22.7	22.6	21.1	20.6	20.8	21.4	21.3	21.4
95% CI Upper	23.4	23.8	23.5	21.8	21.6	21.6	22.2	22.1	22.0
Height	164.7	152.2	158.6	163.1	151.4	157.2	163.5	151.6	157.6
95% CI Lower	164.1	151.7	158.1	162.4	151.0	156.8	163.1	151.2	157.3
95% CI Upper	165.3	152.7	159.3	163.5	151.9	157.2	164.0	152.0	158.0
Weight	62.0	53.3	57.8	56.2	48.2	52.2	57.9	49.7	54.0
95% CI Lower	60.4	52.2	56.4	55.1	47.2	51.4	57.1	49.1	53.1
95% CI Upper	63.5	55.0	59.2	57.3	49.2	53.1	58.8	50.5	54.6
Waist circum.	81.0	73.6	77.2	76.1	70.0	73.1	77.4	71.0	74.2
95% CI Lower	79.4	72.3	76.1	74.8	68.8	72.0	76.4	70.1	73.4
95% CI Upper	82.0	75.0	78.3	77.4	71.2	74.1	78.4	72.0	75.1

centimeters with 77 centimeter for urban respondents and 73 centimeters for rural respondents. The waist circumference for male and female was 77 centimeters and 71 centimeters respectively.

Table 5.3 presents the percentage of respondents according to their BMI category and central obesity by sex and the place of residence. In the survey, we find that 25% respondents were under-weight which was 17%

Table 5.3 Percentage of respondents according to BMI categories by sex and place of residence, Andhra Pradesh, 2007- 08

Category of BMI	Residence						Combined		
	Urban			Rural					
	Male	Female	Total	Male	Female	Total	Male	Female	Total
Under weight(<18.5)	16.9	17.7	17.3	25.9	29.6	27.7	23.3	26.3	24.8
95% CI Lower	13.6	14.7	14.5	22.2	25.8	24.6	20.5	23.4	22.4
95% CI Upper	20.8	21.1	20.4	30.0	33.7	31.1	26.4	29.4	27.3
Normal weight (18.5-24.9)	53.2	51.0	52.2	60.2	54.3	57.3	58.2	53.4	55.9
95% CI Lower	49.9	47.6	49.6	56.4	51.1	54.3	55.3	50.9	53.6
95% CI Upper	56.5	54.5	54.8	64.0	57.5	60.2	61.1	55.9	58.1
Grade-1 over weight (25.0-29.9)	23.6	21.4	22.5	11.6	12.5	12.0	15.1	15.0	15.0
95% CI Lower	20.3	18.6	19.9	9.4	10.1	10.2	13.1	13.0	13.4
95% CI Upper	27.2	24.4	25.4	14.3	15.5	14.2	17.2	17.2	16.8
Grade-2 over weight (30.0-39.9)	6.0	9.5	7.7	1.8	3.5	2.7	3.0	5.2	4.1
95% CI Lower	4.4	7.6	6.2	1.0	2.5	1.9	2.2	4.2	3.4
95% CI Upper	8.3	11.7	9.4	3.4	5.0	3.7	4.2	6.4	4.9
Grade-3 over weight (≥ 40.0)	0.3	0.4	0.3	0.4	0.1	0.2	0.4	0.2	0.3
95% CI Lower	0.1	0.2	0.2	0.1	0.0	0.1	0.1	0.1	0.1
95% CI Upper	0.7	0.8	0.6	1.4	0.3	0.7	1.0	0.3	0.5
Central Obesity WC \geq K*	24.8	36.0	30.3	12.7	21.2	17.0	16.2	25.4	20.8
95% CI Lower	21.7	32.3	27.4	9.8	18.0	14.4	13.8	22.7	18.6
95% CI Upper	28.2	39.9	33.3	16.4	24.8	19.9	19.0	28.2	23.1

* K=90 cm for male and K=80 cm for female

for urban and 28% for rural respondents. By sex, 23% males and 26% females were underweight. Overall 19% population was over weight (30% for urban and 17% for rural respondents). Overall the central obesity was 21% (25% among urban males, 36% among urban females; 13% among rural males, 18% among rural females).

5.5 SOCIO-DEMOGRAPHIC PATTERN OF OVERWIGHT

Overweight (obesity) is a major risk factor of NCD. The socio-demographic pattern of respondents in the category of overweight (grade I, II & III) across age, education, occupation and sex are presented in Table 5.4. Among the urban population, the prevalence of overweight was 31% and the pattern was recorded as increasing with age (9% in 15-24, 33% in 25-34, 46% in 35-44 and 45-54 and 43% in 55-64). The prevalence

among female respondents was high (31%) compare with males (30%), but the increasing pattern with age was observed in both sexes (Table 5.4). In educational categories, the prevalence was varying with 24% among illiterate to 40% among higher level. Occupational categories, the prevalence was high among executive and business class (45%) followed by domestic and service class (36%). Low prevalence of overweight was recorded among agriculture (19%). Overall prevalence among rural population was 15%, which was varying with age (7% in 15-24 to 20% in 55-64). Similarly, prevalence in the educational categories was varying from 13% among illiterate to 20% among higher level of education. In the occupational category, the prevalence was high among executive and business class (28%), and service and sales (26%) and it was low among the people working in agriculture (11%). Overall, prevalence of overweight was

Table 5.4 Percentage of respondents in the category of overweight (Grade I, II & III) across age, education, occupation and by sex and residence, Andhra Pradesh, 2007- 08

Characteristic	Overweight (Grade I, II & III)								
	Urban			Rural			Combined		
	Male	Female	Total	Male	Female	Total	Male	Female	Total
Age group									
15-24	9.0	9.6	9.3	3.2	6.6	4.8	5.0	7.5	6.2
25-34	35.8	29.5	32.7	17.6	17.0	17.3	23.0	20.6	21.8
35-44	43.4	48.5	45.8	20.9	22.4	21.6	27.3	29.6	28.4
45-54	42.3	49.8	45.7	17.8	21.0	19.4	24.6	28.4	26.4
55-64	38.8	47.7	43.3	15.8	20.3	18.2	21.4	26.5	24.1
Total	29.9	31.3	30.5	13.8	16.1	15.0	18.5	20.3	19.4
Education									
Illiterate	16.6	27.2	24.1	10.3	14.3	12.7	11.1	16.4	14.5
Primary	27.6	35.3	31.7	10.7	24.3	16.8	14.0	27.0	20.1
Middle	23.5	34.4	28.7	14.5	19.2	16.5	17.1	24.2	20.3
Secondary	29.4	31.3	30.3	18.7	15.7	17.7	22.1	22.3	22.2
Higher Secondary	28.1	29.1	28.5	15.5	6.5	13.0	21.0	18.7	20.2
College & above	42.6	35.6	40.5	21.5	12.9	19.7	33.5	28.3	32.1
Total	29.9	31.3	30.5	13.8	16.1	15.0	18.5	20.3	19.4
Occupation									
Executive/Business	46.5	34.9	44.5	27.9	28.6	28.1	39.0	31.3	37.3
Agriculture	19.4	18.0	18.8	11.1	11.5	11.3	11.4	11.8	11.6
Domestic Work	**	36.2	36.0	**	20.2	20.5	**	26.3	26.3
Services/Sales	38.6	27.7	36.0	31.9	8.2	26.3	34.9	17.1	30.7
Manual Worker	20.7	14.3	19.5	12.6	16.9	13.9	15.1	16.4	15.5
Other	19.1	16.8	18.4	9.1	7.5	8.6	12.8	11.1	12.3
Total	29.9	31.3	30.5	13.8	16.1	15.0	18.5	20.3	19.4
Number (n)	1257	1459	2716	1462	1978	3440	2719	3437	6156

** Figure not shown; based on fewer than 15 unweighted cases

19% and pattern of prevalence was increasing with age. Except the younger age group, the overweight people are prevalent in all age groups, educational levels and occupation.

In the category of BMI and central obesity, one in

every ten adults in survey population was overweight or categorized into central obesity, constituting a high-risk group for NCD. It is also to be noted that one-fourth of adult population was recorded as under weight which is also an important issue for health planner.



CHAPTER 6

Summary and Conclusions

The NCD risk factors survey in Andhra Pradesh collected information from a random sample of 4905 households covering 2431 households from rural area and 2474 from urban area. From these households, 6218 individuals selected randomly were interviewed to collect behavioural information and also to carry out physical measurements. The analysis of the survey data have been presented and discussed in the present report providing information about the proportion of population or subgroup of population under the risk of non-communicable diseases.

Majority households (82%) in Andhra Pradesh are Hindu followed by Christian (11%). Three-fourth households (94% in urban area and 68% in rural area) had access to piped drinking water. In every ten households, three households in rural area and nine households in urban area had flush toilet facility. Ninety three percent households (91% in rural area and 99% in urban area) in the state used electricity as main source of lighting. Wood was used as a main source of cooking fuel by 81% of rural households whereas 72% of urban households were using LPG. One-fourth of rural households in Andhra Pradesh resided in *kachha* houses and about 55% population was literate. However, there existed sex and rural-urban differentials in educational attainment in the state.

Tobacco is one of the major risk factors of non-communicable diseases. One-third males and very small percentage of females in Andhra Pradesh were found to have smoked tobacco daily. On the other hand, 9% of population (13% of men and 5% of women) used smokeless tobacco. Tobacco in any form - smoking or smokeless used by twenty four percent of population in Andhra Pradesh. This prevalence was 39% among males and 8% among females. The mean age of initiation of tobacco use among male adults age 15-34 was 19 years for smokers and 20 years for smokeless tobacco users.

The alcohol consumption is a known risk factor of many non-communicable diseases. The survey found that about 37% of men consumed alcohol at least once in last one year and 27% of men in last one month. The alcohol

consumption among females was very low. Those who consumed alcohol in last seven days, 23% of them were binge drinkers. The mean age of initiation of alcohol consumption among men age 15-34 who ever consumed alcohol was 20 years.

Nutritional inadequacy is the major risk factor of many non-communicable diseases. In Andhra Pradesh, 88% population consumed less than five servings of fruits and vegetables per day, which was inadequate as per WHO recommended standards. On an average, people consumed fruits only two days in a week against vegetables, which was consumed 5 days in a week.

Physical inactivity is the leading cause of diabetes, hypertension and coronary heart diseases. In Andhra Pradesh, 68% of population (78% of urban and 64% of rural population) was in low category of physical activity. When tested for hypertension, 43% of the population was detected with pre-hypertension stage and one-fifth was in stage I and stage II hypertension. As per BMI status, 13% of population was in the category of over weight and one-quarter population was recorded as under weight. However, 14% of population (17% among females and 11% among males) in Andhra Pradesh was in the category of central obesity.

Overall, prevalence of smoking and smokeless tobacco use among female population was low compare with males and increasing pattern of prevalence was recorded with increasing age group of people. A declining pattern of prevalence was observed with increasing level of education. Prevalence among the people working in agriculture and as manual worker was high compare with others. A similar pattern of increasing prevalence with age and decreasing with level of education was also observed with current alcohol drinkers. The habits of tobacco and alcohol use starts at early young age which contributes to the high risk of NCD at later age. High proportion of population was taking inadequate amount of fruits and vegetables which increases the risk of NCD. Its distribution across all age groups, education and occupation by sex and residence was found high with marginal differences. Besides that, more than half of the population was found in the category of doing low

physical activity. More female respondents were in the category of low physical activity as compare with males across all the age groups. Rural population was doing more physical work than urban. The increasing pattern of prevalence of hypertension was recorded with increasing age group of people. It was prevalent in all education levels and occupational categories. High prevalence of overweight was recorded in all the age groups except the younger age. It was prevalent in both

sexes, but higher in urban population compare with rural. Low prevalence of overweight was recorded among illiterates as well as among the people working in agriculture or manual worker. Overall, NCD risk factors were prevalent across all the socioeconomic and demographic categories of population in Andhra Pradesh.

These are the major health issues related to non-communicable diseases of people in Andhra Pradesh.



References

1. Risk factor surveillance for noncommunicable diseases (NCDs): the multi-site ICMR-WHO collaborative initiative, presented in Forum 9, Mumbai, 12-16 September 2005.
2. The World Health Report 2002; Reducing Risks, Promoting Healthy Life, WHO.
3. Primary Census Abstract: Census of India, 2001, Office of the Registrar General, India, Ministry of Home Affairs, Govt. of India.
4. National Health Profile, Central Bureau of Health Intelligence, Director General Health Services, Ministry of Health & Family Welfare, Govt. of India.
5. Lisle Kish: A procedure for objective respondent selection within the household. *Am Stat Assoc Journal*, 1949; 44: 380-387.
6. Lehtonen R. and Pahkinen E. 2004. *Practical Methods for Design and Analysis of Complex Surveys*. John Wiley & Sons, Ltd.: England.
7. Global Physical Activity Questionnaire (GPAQ), Analysis Guide, www.who.int/chp/steps.
8. The Seventh Report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure. *JAMA* 2003;289:2560-71.
9. WHO/IASO/IOTF: The Asia-Pacific perspective: redefining obesity and its treatment [article online]. Available from <http://www.idi.org.au/home.htm>. Accessed 21 August 2003.



Appendix - A

SAMPLE WEIGHTS

First, appropriate sampling weights for households were constructed for each state data set separately for Urban and Rural sectors. The element weight consisted of factors reflecting ward selection probabilities, Census enumeration block (CEB) selection probabilities within wards; and household selection probabilities within CEB; and household non-response adjustments.

For Urban area of a state, the weight HWT_{ijk} for the household k in CEB j of ward i , can be expressed as follows

$$HWT_{ijk} = w_{1i} \times w_{2ji} \times w_{3k|i,j} \quad i=1, \dots, 50, j=1, \dots, 50$$

where $w_{1i} = \frac{1}{\pi_i}$: the reciprocal of the inclusion probability π_i of ward i

$$\text{where } \pi_i = \frac{a \times \text{Population of ward } i}{\text{Total Urban Population}} \quad \text{and}$$

$a (=50)$ is the total number wards to be selected from the urban sector

$w_{2ji} = \frac{1}{\pi_{j|i}}$: the reciprocal of the conditional probability of selection of CEB j in ward i

$$\text{where } \pi_{j|i} = \frac{\text{Population of selected CEB } j \text{ within ward } i}{\text{Population of selected ward } i}$$

$w_{3k|i,j} = \frac{1}{\pi_{k|i,j} \times \hat{\theta}_{k|i,j}}$: the reciprocal of the product of conditional inclusion probability $\pi_{k|i,j}$ of household k in the j^{th} selected CEB of the i^{th} ward and estimated conditional response probability $\hat{\theta}_{k|i,j}$ of household k from within the j^{th} selected CEB of ward i .

$$\text{where } \pi_{k|i,j} = \frac{\text{Number of households sampled from selected CEB } j \text{ of ward } i}{\text{Number of households in selected CEB } j \text{ of ward } i}$$

$$HWT_{ijk} = \frac{\text{Size of Urban Population}}{50 \times \text{Population of selected CEB from ward } i} \times \frac{\text{Number of households in selected CEB of ward } i}{\text{Number of households sampled from selected CEB of ward } i \text{ with HH Result code completed}}$$

In rural sector, from the lists of villages, 50 villages (or cluster of villages) were selected with probability proportional to size and from each village 50 household were selected using systematic sampling.

Proceeding as above it can be shown the weight for the k^{th} selected household of the i^{th} selected village, HWT_{ijk} ,

$$HWT_{ijk} = \frac{\text{Size of Rural Population}}{50 \times \text{Population of } i^{\text{th}} \text{ selected village}} \times \frac{\text{Number of households in } i^{\text{th}} \text{ selected village}}{\text{Number of households selected from } i^{\text{th}} \text{ village with HH Result code complete}}$$

INDIVIDUAL WEIGHTS

From each selected household one member aged 15-54 is selected using the Kish Method and all usual members aged 55-64 were selected. Since objective of the study is to obtain estimates for each age group (15-24 through 55-64) and sex groups, post stratification is used for improvement of efficiency of the estimators.

Post stratification weights for individuals were constructed using the state age distributions for both sexes of the urban sector which are available on the population level. We first divide the target population of persons age 15-64 in 10 age - sex post strata with five age group (15-24 through 55-64) and two sex groups (male and female).

In the subsequent lines the symbol l is used to denote the age group $[15 + (l - 1)*10, 15 + 10*l]$, $l = 1, 2, \dots, 5$ and m for sex, $m = 1$ if sex is male and $m=2$ if sex is female.

For Urban,

Define :

$$\delta_{ijknlm} = \begin{cases} 1 & \text{if } n^{\text{th}} \text{ selected respondent of the } k^{\text{th}} \text{ household of the } j^{\text{th}} \text{ CEB of the } i^{\text{th}} \\ & \text{ward belongs to age group } l \text{ and of sex } m. \\ 0 & \text{otherwise} \end{cases}$$

$$\hat{N}_{lm} \begin{cases} \text{estimated number of persons of age group } l \text{ and sex } m \text{ if one person from the list of persons age} \\ \text{15-54 is selected from household of the population } (l = 1, 2, 3, 4, m = 1, 2) \\ \text{estimated number of the persons belonging to the age group } l \text{ and sex group } m (l = 5, m = 1, 2) \end{cases}$$

\hat{N}_{lm} is obtained as

$$\hat{N}_{lm} = \frac{1}{\hat{\theta}_{lm}} \sum_{\substack{\text{over all} \\ \text{all possible} \\ \text{values of} \\ i, j, k, n}} HWT_{ijk} \times \delta_{ijknlm} \quad \text{where } \hat{\theta}_{lm} \text{ is the estimated group response rate.}$$

Calibrated Individual weight

$$IWT_{ijklm} = \frac{N_{lm}}{\hat{N}_{lm}} \times HWT_{ijk}$$

Denoted by

N_{lm} = Number of person of sex m belonging to age group l in the urban sector of the population
($l = 1, 2, 3, 4, 5$ and sex $m = 1, 2$)

y_{ijkn} = the observed value of the study variable for the respondent n belonging to household k ,
CEB j and ward i .

Estimate of the population total of sex group m and age group l is

$$\hat{Y}_{lm} = \sum_{\substack{\text{over all} \\ \text{all possible} \\ \text{values of} \\ i, j, k, n}} \delta_{ijknlm} \times IWT_{ijklm}$$

$$\hat{N}_l = \hat{N}_{l1} + \hat{N}_{l2} \quad , \quad \hat{Y}_l = \hat{Y}_{l1} + \hat{Y}_{l2} \quad , \quad l = 1, \dots, 5$$

$$\hat{N}_m = \hat{N}_{1m} + \dots + \hat{N}_{5m} \quad , \quad \hat{Y}_m = \hat{Y}_{1m} + \dots + \hat{Y}_{5m} \quad , \quad m = 1, 2$$

$$\hat{N} = \sum_{l=1}^5 \sum_{m=1}^2 N_{l,m} \quad , \quad \hat{Y} = \sum_{l=1}^5 \sum_{m=1}^2 \hat{Y}_{l,m}$$

Estimate of the mean of the study variable for sex group m and age group l, \widehat{Y}_{lm} and for and overall are $\frac{\widehat{Y}_{lm}}{\widehat{N}_{lm}}, \frac{\widehat{Y}_m}{\widehat{N}_m}, \frac{\widehat{Y}_l}{\widehat{N}_l}, \frac{\widehat{Y}}{\widehat{N}}$, respectively.

For Rural,

Define :

$$\delta_{iknlm} = \begin{cases} 1 & \text{if } n^{\text{th}} \text{ selected respondent of the } k^{\text{th}} \text{ household of the } i^{\text{th}} \text{ village} \\ & \text{belongs to age group l and of sex m.} \\ 0 & \text{otherwise} \end{cases}$$

$$\widehat{N}_{lm} = \begin{cases} \text{estimated number of persons of age group l and sex m if one person from the list of} \\ \text{persons age 15-54 is selected from household of the population (l =1,2,3,4, m=1,2)} \\ \text{estimated number of the persons belonging to the age group l and sex group m (l =5, m=1,2)} \end{cases}$$

\widehat{N}_{lm} is obtained as

$$\widehat{N}_{lm} = \frac{1}{\widehat{\theta}_{l,m}} \sum_{\substack{\text{over all} \\ \text{all possible} \\ \text{values of} \\ i,j,k,n}} HWT_{ik} \times \delta_{ijknlm}, \quad \text{where } \widehat{\theta}_{l,m} \text{ is the estimated group response rate.}$$

Calibrated Individual weight

$$IWT_{iklm} = \frac{N_{lm}}{\widehat{N}_{lm}} \times HWT_{ik}$$

Denoted by

N_{lm} = Number of person of sex m belonging to age group l in the rural sector of the population (l = 1,2 ,3,4,5 and sex m =1,2)

y_{ikn} , = the observed value of the study variable for the respondent n belonging to household k of village i.

Estimate of the population total of sex group m and age group l is

$$\widehat{Y}_{l,m} = \sum_{\substack{\text{over all} \\ \text{all possible} \\ \text{values of} \\ i,j,k,n}} \delta_{ijknlm} \times y_{ikn} \times IWT_{ijkml}$$

Estimate of the mean of the study variable for age-sex group l and m, sex group m, age group l and overall mean can be obtained.

Appendix - B

INTEGRATED DISEASE SURVEILLANCE PROJECT (IDSP)

NCD RISK FACTORS SURVEY (PHASE -I), INDIA

(Name of State Year -2007)

HOUSEHOLD QUESTIONNAIRE

IDENTIFICATION	
STATE :	[] []
DISTRICT :	[] []
TEHSIL/TALUK	[] [] [] []
CITY/TOWN/VILLAGE:	[] [] [] [] [] [] [] []
URBAN/RURAL (<i>URBAN=1, RURAL =2, URBAN SLUM -3</i>)	[]
PSU NUMBER	[] [] []
SEGMENT NUMBER:	[]
Household Number	[] [] [] []
Name of Household Head: _____	
Address of HOUSEHOLD: _____	

INTERVIEWER VISITS				
	1	2	3	FINAL VISIT
Date				Day [] [] Month [] [] Year 2 0 [] []
Interviewer's Name				Interviewer Code [] []
Result				Result* []
Next Visit: Date Time				Total Number of Visits []
<p>*RESULT CODES:</p> <div style="display: flex; justify-content: space-between;"> <div style="width: 48%;"> <p>1. COMPLETED</p> <p>2. NO HOUSEHOLD MEMBER/ NO COMPETENT RESPONDENT AT HOME AT THE TIME OF VISIT</p> <p>3. ENTIRE HOUSEHOLD ABSENT FOR EXTENDED PERIOD</p> <p>4. POSTPONED</p> <p>5. REFUSED</p> </div> <div style="width: 48%;"> <p>6. DWELLING VACANT OR ADDRESS NOT A DWELLING</p> <p>7. DWELLING DESTROYED</p> <p>8. DWELLING NOT FOUND</p> <p>9. OTHER _____ (SPECIFY)</p> </div> </div>				
NAME DATE	SUPERVISOR [] []	EDITED & CHECKED BY [] []	KEYED BY [] []	

HOUSEHOLD STRUCTURE (HS)						
List of all household members who usually live in your household aged 12 years and above						
LINE NO.	NAME	RELATIONSHIP	SEX	AGE IN COMPLETED YEARS	RESIDENTIAL STATUS	RECRUITED FOR SURVEY
	Please give me names of the persons who usually live in your household (may be temporarily away from home)	(With head of household)	Male-1 Female-2		(Present-1; temporarily away from home-2)	Put a tick mark against one member age 15-54 selected below by kish method and all members age 55-64
(1)	(2)	(3)	(4)	(5)	(6)	(7)
01						
02						
03						
04						
05						
06						
07						
08						
09						
10						
11						
12						
Codes for Q.3 Relationship to Head of Household:		01 - HEAD, 02 - WIFE OR HUSBAND, 03 - SON OR DAUGHTER 04 - SON IN LAW OR DAUGHTER IN LAW 05 - GRANDCHILD 06 - PARENT 07 - PARENT IN LAW			08 - BROTHER OR SISTER 09 - BROTHER IN LAW OR SISTER IN LAW 10 - NIECE OR NEPHEW 11 - OTHER RELATIVE 12 - ADOPTED OR FOSTER CHILD 13 - NOT RELATED	

LIST ALL USUAL MEMBERS OF THE HOUSEHOLD AGE 15 - 54 IN THE HOUSEHOLD*.

Line No.	Sex	Age	Adult Number	Select one member (R) by using Kish Table	Enter a specific Kish Table used for selection of one member below. (A or B1 or B2 or C..... as assigned for each randomly selected household 1 to 50)

*Arrange all the members aged 15-54 in the following order - oldest male, next oldest male, and so on for all males followed by oldest female, next oldest female, etc. Then use selection table assigned to the household to choose R individual RESPONDENT.

GENERAL HOUSEHOLD INFORMATION			
Questions		RESPONSE	SKIP
1.	Number of members who usually live in the household	<input type="text"/> <input type="text"/>	
2.	Religion of the head of the household:	Hindu 01 Muslim 02 Christian 03 Sikh 04 Buddhist/neo buddhist 05 Jain 06 Jewish 07 Parsi 08 No religion 09 Other 96 (Specify)	
3.	What is the main source of drinking water?	PIPED WATER Piped into Residence 11 Public Tap 12 GROUND WATER: Hand Pump in Residence 21 Public Hand Pump 22 WELL WATER Well in Residence Covered well 31 Open well 32 Public Well Covered well 33 Open well 34 SURFACE WATER: Spring 41 River/Stream 42 Pond 43 Dam 44 Rainwater 51 Tanker Truck 61 Any other 96 (specify)	
4.	What kind of toilet facilities do you have?	Flush Toilet Own Flush Toilet 11 Shared Flush Toilet 12 Public Flush Toilet 13 Pit Toilet/Latrine Own Pit Toilet 21 Shared Pit Toilet 22 Public Pit Toilet 23 No facility/Bush/Field 31 Other 96 (Specify)	

5.	What is the main source of lighting for your household?	Electricity 1 Kerosene 2 Gas 3 Oil 4 Other 6 (Specify)																																																													
6.	What is the type of house ?	Pucca 1 Semi-Pucca 2 Kachha 3																																																													
7.	How many rooms are there in your household?	Rooms..... <input type="text"/> <input type="text"/>																																																													
8.	Do you have a separate room, which is used as kitchen ?	Yes 1 No 2																																																													
9.	What type of fuel does your household mainly use for cooking?	Wood 01 Crop Residue 02 Dung Cakes 03 Coal/Coke/Lignite 04 Charcoal 05 Kerosene 06 Electricity 07 Liquid Petroleum Gas (LPG) 08 Bio-Gas 09 Others 96 (Specify)																																																													
10.	Does this household own this house or any other house?	Yes 1 No 2																																																													
11.	Does this household own any agriculture land ?	Yes 1 No 2	If No, go to 14																																																												
12.	How much agriculture land does this household own?	Acres <input type="text"/> <input type="text"/> . <input type="text"/> None																																																													
13.	Out of this land, how much is irrigated ?	Acres <input type="text"/> <input type="text"/> . <input type="text"/> None																																																													
14.	Does the household own any livestock?	Yes 1 No 2																																																													
15.	Does the household own any of the following: (<i>READ ALL THE OPTIONS AND RECORD THE RESPONSE</i>)	<table border="0"> <thead> <tr> <th></th> <th>Yes</th> <th>No</th> </tr> </thead> <tbody> <tr> <td>A mattress?</td> <td>1</td> <td>2</td> </tr> <tr> <td>A pressure cooker?</td> <td>1</td> <td>2</td> </tr> <tr> <td>A chair?</td> <td>1</td> <td>2</td> </tr> <tr> <td>A cot or bed?</td> <td>1</td> <td>2</td> </tr> <tr> <td>A table?</td> <td>1</td> <td>2</td> </tr> <tr> <td>A clock or Watch?</td> <td>1</td> <td>2</td> </tr> <tr> <td>An electric fan?</td> <td>1</td> <td>2</td> </tr> <tr> <td>A bicycle?</td> <td>1</td> <td>2</td> </tr> <tr> <td>A radio or transistor?</td> <td>1</td> <td>2</td> </tr> <tr> <td>A sewing machine?</td> <td>1</td> <td>2</td> </tr> <tr> <td>A telephone or Mobile?</td> <td>1</td> <td>2</td> </tr> <tr> <td>A refrigerator?</td> <td>1</td> <td>2</td> </tr> <tr> <td>A television?</td> <td>1</td> <td>2</td> </tr> <tr> <td>A moped, scooter, or motorcycle?</td> <td>1</td> <td>2</td> </tr> <tr> <td>A car?</td> <td>1</td> <td>2</td> </tr> <tr> <td>A water pump?</td> <td>1</td> <td>2</td> </tr> <tr> <td>A bullock cart?</td> <td>1</td> <td>2</td> </tr> <tr> <td>A thresher?</td> <td>1</td> <td>2</td> </tr> <tr> <td>A tractor?</td> <td>1</td> <td>2</td> </tr> </tbody> </table>		Yes	No	A mattress?	1	2	A pressure cooker?	1	2	A chair?	1	2	A cot or bed?	1	2	A table?	1	2	A clock or Watch?	1	2	An electric fan?	1	2	A bicycle?	1	2	A radio or transistor?	1	2	A sewing machine?	1	2	A telephone or Mobile?	1	2	A refrigerator?	1	2	A television?	1	2	A moped, scooter, or motorcycle?	1	2	A car?	1	2	A water pump?	1	2	A bullock cart?	1	2	A thresher?	1	2	A tractor?	1	2	
	Yes	No																																																													
A mattress?	1	2																																																													
A pressure cooker?	1	2																																																													
A chair?	1	2																																																													
A cot or bed?	1	2																																																													
A table?	1	2																																																													
A clock or Watch?	1	2																																																													
An electric fan?	1	2																																																													
A bicycle?	1	2																																																													
A radio or transistor?	1	2																																																													
A sewing machine?	1	2																																																													
A telephone or Mobile?	1	2																																																													
A refrigerator?	1	2																																																													
A television?	1	2																																																													
A moped, scooter, or motorcycle?	1	2																																																													
A car?	1	2																																																													
A water pump?	1	2																																																													
A bullock cart?	1	2																																																													
A thresher?	1	2																																																													
A tractor?	1	2																																																													

Integrated Disease Surveillance Project (IDSP)

16.	<p>What is the type of oil/cooking medium most commonly used in the house?</p> <p>(CHOOSE ONLY ONE IDENTIFIED BY MAXIMUM CONSUMPTION)</p>	<p>Cooking Oil</p> <p>Mustard oil 01</p> <p>Coconut oil 02</p> <p>Groundnut oil 03</p> <p>Sunflower oil 04</p> <p>Soyabean oil 05</p> <p>Palm oil 06</p> <p>Vanaspati oil 07</p> <p>Pure Ghee 08</p> <p>Butter 09</p> <p>Others 96</p> <p>(Specify)</p>	
-----	--	---	--

STEP- I DEMOGRAPHIC INFORMATION																																							
QUESTIONS AND FILTERS		Response	Skip																																				
101.	Sex	Male 1 Female 2																																					
102.	Age	Age in completed Years <input type="text"/> <input type="text"/>																																					
103.	What is your current marital status ?	Never married 1 Currently 2 Married 2 Married but gauna not performed..... 3 Widowed/Divorced/Separated 4																																					
104.	Have you ever attended school?	Yes 1 No 2	If no, go to 107																																				
105.	if yes , what is the highest grade of education you completed?	Grade*..... <input type="text"/> <input type="text"/>																																					
106.	Check 105 Grade 0-5 <input type="text"/> ↓	Grade 6 & above <input type="text"/> → Go to 108																																					
107.	Can you read and write?	Yes 1 No 2																																					
108.	What is your main work/ occupation?	Professional/Executive/Manager/ Big business 1 Clerical/Medium business 2 Sales 3 Agriculture/Self-employed 4 Agriculture employer 5 Household and domestic work 6 Services 7 Skilled manual 8 Unskilled manual 9 Other (Specify)..... 10 Do not work 11																																					
<p>*GRADE FOR DIFFERENT LEVEL OF COMPLETED EDUCATION</p> <table border="0"> <thead> <tr> <th></th> <th>EDUCATION LEVEL</th> <th></th> <th>GRADE</th> </tr> </thead> <tbody> <tr> <td></td> <td>CLASS I TO XII</td> <td>: 1 TO 12 YEARS</td> <td>= 1 TO 12 GRADE</td> </tr> <tr> <td></td> <td>BACHELOR'S DEGREE</td> <td>: 15 YEARS (12+3)</td> <td>= 15 GRADE</td> </tr> <tr> <td></td> <td>MASTER'S DEGREE</td> <td>: 17 YEARS (12+3+2)</td> <td>= 17 GRADE</td> </tr> <tr> <td></td> <td>ENGINEERING</td> <td>: 16 YEARS (12+4)</td> <td>= 16 GRADE</td> </tr> <tr> <td></td> <td>MBBS</td> <td>: 17 YEARS (12+5)</td> <td>= 17 GRADE</td> </tr> <tr> <td></td> <td>POLYTECHNIC</td> <td>: 13 YEARS (10+3)</td> <td>= 13 GRADE</td> </tr> <tr> <td></td> <td>ITI</td> <td>: 11 YEARS (10 +1)</td> <td>= 11 GRADE</td> </tr> <tr> <td></td> <td>PH. D.</td> <td>: 20 YEARS (12+3+2+3)</td> <td>= 20 GRADE</td> </tr> </tbody> </table>					EDUCATION LEVEL		GRADE		CLASS I TO XII	: 1 TO 12 YEARS	= 1 TO 12 GRADE		BACHELOR'S DEGREE	: 15 YEARS (12+3)	= 15 GRADE		MASTER'S DEGREE	: 17 YEARS (12+3+2)	= 17 GRADE		ENGINEERING	: 16 YEARS (12+4)	= 16 GRADE		MBBS	: 17 YEARS (12+5)	= 17 GRADE		POLYTECHNIC	: 13 YEARS (10+3)	= 13 GRADE		ITI	: 11 YEARS (10 +1)	= 11 GRADE		PH. D.	: 20 YEARS (12+3+2+3)	= 20 GRADE
	EDUCATION LEVEL		GRADE																																				
	CLASS I TO XII	: 1 TO 12 YEARS	= 1 TO 12 GRADE																																				
	BACHELOR'S DEGREE	: 15 YEARS (12+3)	= 15 GRADE																																				
	MASTER'S DEGREE	: 17 YEARS (12+3+2)	= 17 GRADE																																				
	ENGINEERING	: 16 YEARS (12+4)	= 16 GRADE																																				
	MBBS	: 17 YEARS (12+5)	= 17 GRADE																																				
	POLYTECHNIC	: 13 YEARS (10+3)	= 13 GRADE																																				
	ITI	: 11 YEARS (10 +1)	= 11 GRADE																																				
	PH. D.	: 20 YEARS (12+3+2+3)	= 20 GRADE																																				

STEP- I BEHAVIOURAL INFORMATION

Now I am going to ask you some questions about various health behaviours. This includes things like smoking, drinking alcohol, eating fruits and vegetables and physical activity. Let's start with tobacco

Smoking Tobacco use:

Questions		Response	Skip														
201.	Do you currently smoke any tobacco products, such as bidis, cigarettes, cigars or pipes, hookah or any other local tobacco products?	Yes 1 No..... 2	If No, go to 205														
202.	If Yes , do you smoke daily ?	Yes 1 No..... 2	if No, go to 205														
203.	On an average, how many (number of times in case of hookah) of the following do you smoke each day? <i>(RECORD FOR EACH TYPE)</i> <i>RECORD 88, IF ANY PRODUCT IS NOT USED INSTEAD OF LEAVING BLANK IN THE PRODUCT CATEGORIES).</i> <i>(RECORD FOR ANY NEW FORM OF TOBACCO USE REPORTED BY THE RESPONDENT e.g. REVERSE SMOKING etc.)</i>	<table border="0" style="width: 100%;"> <tr> <td style="text-align: right;">Bidis</td> <td style="text-align: center;">Number <input type="text"/> <input type="text"/></td> </tr> <tr> <td style="text-align: right;">Manufactured Cigarettes</td> <td style="text-align: center;"><input type="text"/> <input type="text"/></td> </tr> <tr> <td style="text-align: right;">Hand-rolled Cigarettes</td> <td style="text-align: center;"><input type="text"/> <input type="text"/></td> </tr> <tr> <td style="text-align: right;">Pipes</td> <td style="text-align: center;"><input type="text"/> <input type="text"/></td> </tr> <tr> <td style="text-align: right;">Cigars, Cheroots</td> <td style="text-align: center;"><input type="text"/> <input type="text"/></td> </tr> <tr> <td style="text-align: right;">Hookah</td> <td style="text-align: center;"><input type="text"/> <input type="text"/></td> </tr> <tr> <td style="text-align: right;">Other local smoked tobacco products..... (SPECIFY)</td> <td style="text-align: center;"><input type="text"/> <input type="text"/></td> </tr> </table>	Bidis	Number <input type="text"/> <input type="text"/>	Manufactured Cigarettes	<input type="text"/> <input type="text"/>	Hand-rolled Cigarettes	<input type="text"/> <input type="text"/>	Pipes	<input type="text"/> <input type="text"/>	Cigars, Cheroots	<input type="text"/> <input type="text"/>	Hookah	<input type="text"/> <input type="text"/>	Other local smoked tobacco products..... (SPECIFY)	<input type="text"/> <input type="text"/>	
Bidis	Number <input type="text"/> <input type="text"/>																
Manufactured Cigarettes	<input type="text"/> <input type="text"/>																
Hand-rolled Cigarettes	<input type="text"/> <input type="text"/>																
Pipes	<input type="text"/> <input type="text"/>																
Cigars, Cheroots	<input type="text"/> <input type="text"/>																
Hookah	<input type="text"/> <input type="text"/>																
Other local smoked tobacco products..... (SPECIFY)	<input type="text"/> <input type="text"/>																
204.	How old were you at that time when you first started using the tobacco product(s) daily ?	Age in completed years <input type="text"/> <input type="text"/> Don't remember 7 7	Go to 208														
205.	In the past, did you ever smoke tobacco products such as bidis, cigarettes, cigars or pipes daily ?	Yes 1 No 2	If No, go to 207														
206.	How old were you when you stopped smoking daily?	Age in completed years <input type="text"/> <input type="text"/> Don't remember 7 7															
207.	Are you currently exposed to tobacco smoke at your home or workplace daily ?	Yes 1 No 2															

Smokeless Tobacco use			
Questions		Response	Skip
208.	Do you currently use any smokeless tobacco , such as (chewing tobacco, <i>tuibu</i> snuff, betel, gutka, pan masala, etc.)?	Yes 1 No 2	if No, go to 212
209.	If yes , Do you currently use smokeless tobacco products daily ?	Ye 1 No 2	if No, go to 212
210.	On average, how many times a day do you use... <i>(RECORD FOR EACH TYPE)</i> <i>SPECIFY 77 IF NO PRODUCTS WERE USED IN EACH CATEGORY INSTEAD OF LEAVING CATEGORIES BLANK.</i>	Chewing tobacco <input type="text"/> <input type="text"/> Pan with tobacco <input type="text"/> <input type="text"/> Tuibu, Tobacco Snuff, by mouth <input type="text"/> <input type="text"/> Snuff, by nose <input type="text"/> <input type="text"/> Other <input type="text"/> <input type="text"/> Other (specify).....	
211.	How old were you at that time when you first started using smokeless tobacco daily ?	Age in completed years <input type="text"/> <input type="text"/>	Go to 214
212.	If you are not using currently, in the past did you ever use smokeless tobacco products daily such as chewing tobacco, tuibu, snuff, betel, gutka, etc.?	Yes 1 No 2	if No, go to 214
213.	How old were you when you stopped using smokeless tobacco products daily ?	Age in completed years <input type="text"/> <input type="text"/>	

Alcohol Consumption			
The next questions ask about the consumption of alcohol.			
Questions		Response	Skip
214.	Have you consumed any alcoholic products (such as beer, wine, whisky, locally prepared alcohol, etc.) within the past 12 months ?	Yes 1 No 2	if No, go to 219
215.	In the past 12 months, how frequently have you had at least one drink?	5-7 days per week 1 1-4 days per week 2 1-3 days per month 3 Less than once per month 4	
216.	When you drink alcohol, on average , how many standard drinks do you have during one day? (<i>USE SHOWCARD</i>)	Number <input type="text"/> <input type="text"/>	
217.	Have you consumed alcohol (such as beer, wine, spirits, or any locally prepared wine, etc.) within the past 30 days ?	Yes 1 No 2	If No go to 220
218.	During each of the past 7 days, how many standard drinks of any alcoholic drink did you have each day? <i>(USE SHOWCARD)</i>	Monday <input type="text"/> <input type="text"/> Tuesday <input type="text"/> <input type="text"/> Wednesday <input type="text"/> <input type="text"/>	Go to 220

		Thursday <input type="text"/>	
		Friday <input type="text"/>	
		Saturday <input type="text"/>	
		Sunday <input type="text"/>	
219.	If answer to Question 214 is No, then Have you ever (past user) consumed alcohol (such as beer, wine, spirits, or any local wine product)?	Yes 1 No 2	if No, go to 221
220.	How old were you when you started consuming alcohol regularly?	Age in years <input type="text"/> Don't Remember 7 7	

Diet

The next questions ask about the fruits and vegetables that you usually eat. I have a nutrition card here that shows you some examples of local fruits and vegetables. Each picture represents the size of a serving. As you answer these questions please think of a 'typical' or a 'usual' week.

Questions		Response	Skip
221.	In a typical week, on how many days do you eat fruit?	Number of days <input type="text"/>	If zero days, go to 223
222.	How many servings of fruit do you eat on one of those days? (USE SHOWCARD)	Number of servings <input type="text"/>	
223.	In a typical week, how many days do you eat vegetables? (USE SHOWCARD)	Number of days <input type="text"/>	If zero days, go to 225
224.	How many servings of vegetables do you eat on one of those days? (USE SHOWCARD)	Number of servings <input type="text"/>	
225.	How often do you consume each of the following ? (USE CODE: DAILY - 1; AT LEAST ONCE IN A WEEK - 2; ONCE IN A MONTH -3; OCCASIONALLY OR RARELY - 4; NEVER - 5)	Butter/Ghee <input type="text"/> Fried local foods (Samosa, Kachori, etc.) <input type="text"/> Red meat <input type="text"/> Eggs <input type="text"/> Chicken <input type="text"/> Fish <input type="text"/> Aerated drinks <input type="text"/> Sweetened drinks <input type="text"/> Pizza/burgers/French fries etc <input type="text"/> Cakes, Pastries or other bakery items <input type="text"/> Chips, Namkeen etc <input type="text"/>	

Physical Activity

Next I am going to ask you about the time you spend doing different types of physical activity in a **typical week**. Please answer these questions even if you do not consider yourself to be a physically active person.

Think first about the time you spend doing work. Work includes things that you have to do such as paid or unpaid work, study/training, household chores, harvesting food/crops, fishing or hunting for food, seeking employment.

In answering the following questions '**Vigorous -Intensity activities**' are activities that require hard physical effort and cause large increase in breathing or heart rate, '**Moderate-Intensity activities**' are activities that require effort and cause small increases in breathing or heart rate.

Questions		Response	Skip
226.	Does your work involve vigorous-intensity activity that causes large increases in breathing or heart rate like (carrying or lifting heavy loads, digging or construction work etc.) for at least 10 minutes continuously ?	Yes 1 No 2	If No, go to 229
227.	In a typical week, on how many days do you do vigorous-intensity activities as part of your work?	Number of days <input type="text"/>	
228.	How much time do you spend doing vigorous-intensity activity at home/work on a typical day?	Hours : minutes <input type="text"/> : <input type="text"/> Hours minutes	
229.	Does your work involve moderate -intensity activity that causes small increases in breathing or heart rate for at least 10 minutes continuously (such as brisk walking or carrying loads, manual washing of clothes, dry sweeping of floor, wet mopping of floor, drawing water from well, carrying water from tap, carrying water from river or well, manual grinding or pounding of cereals, gardening at home, carrying groceries from market, etc.) ?	Yes 1 No 2	If No, go to 232
230.	In a typical week, on how many days do you do moderate-intensity activities as part of your work?	Number of days <input type="text"/>	
231.	How much time do you spend doing moderate-intensity activity at work on a typical day?	Hours : minutes <input type="text"/> : <input type="text"/> Hours minutes	

Travel (related to Physical Activity) to and from places

The next questions exclude the physical activities at work that you have already mentioned. Now I would like to ask you about the usual way you travel to and from places. For example to work, for shopping, to market, to place of worship etc.

232.	Do you walk or use a bicycle (pedal cycle) for at least 10 minutes continuously to get to and from places?	Yes 1 No 2	If No, go to 235
233.	In a typical week, on how many days do you walk or bicycle for at least 10 minutes continuously to get to and from places?	Number of days <input type="text"/>	
234.	How much time do you spend walking or bicycling for travel on a typical day?	Hours : minutes <input type="text"/> : <input type="text"/> Hours minutes	

Recreational Activity			
235.	Do you do any vigorous-intensity sports , fitness or recreational (leisure) activities that cause large increases in breathing or heart rate like (running or football,...) for at least 10 minutes continuously ?	Yes 1 No..... 2	If No, go to 238
236.	In a typical week, on how many days do you do vigorous-intensity sports, fitness, or recreational activity?	Number of days <input type="text"/>	
237.	How much time do you spend doing vigorous-intensity sports, fitness or recreational (leisure) activities on a typical day ?	Hours : minutes <input type="text"/> : <input type="text"/> Hours minutes	
238.	Do you do any moderate-intensity sports , fitness or recreational (leisure) activities that cause small increases in breathing or heart rate such as brisk walking (cycling, swimming, volleyball etc.) for at least 10 minutes continuously ?	Yes 1 No..... 2	If No, go to 241
239.	In atypical week, on how many days do you do moderate-intensity sports, fitness, or recreational activity?	Number of days <input type="text"/>	
240.	How much time do you spend doing moderate-intensity sports, fitness, or recreational activities on a typical day?	Hours : minutes <input type="text"/> : <input type="text"/> Hours minutes	
Yoga Activity			
241.	Do you regularly practice Yogic Exercise /Yogasan ?	Yes 1 No..... 2	If No, go to 244
242.	If yes , how many days in a week?	Number of days <input type="text"/>	
243.	How much time do you spend doing Yoga in a typical day ?	Hours : minutes <input type="text"/> : <input type="text"/> Hours minutes	
Sedentary Behaviour			
The following question is about sitting or reclining at work, at home, getting to and from places, or with friends including time spent [sitting at a desk, sitting with friends, traveling in car, bus, train, reading, playing cards or watching television], but do not include time spent sleeping.			
244.	How much time do you usually spend sitting or reclining on a typical day ?	Hours : minutes <input type="text"/> : <input type="text"/> Hours minutes	

History of Raised Blood Pressure			
Questions		Response	Skip
245.	When was your blood pressure last measured by a health professional ?	Within past 12 months 1 1-5 years ago 2 More than 5 years ago 3 Never 4	
246.	Have you ever been told by a doctor or other health worker that you have raised (high) blood pressure or hypertension?	Yes 1 No 2	If No, go to 248
247.	Are you currently receiving any of the following treatments/advice for raised (high) blood pressure prescribed by a doctor or other health worker as well as any advice ?		
	Drugs (medication) that you have taken in the last 2 weeks	Yes 1 No 2	
	Special prescribed diet	Yes 1 No 2	
	Advice or treatment to lose weight	Yes 1 No 2	
	Advice or treatment to stop smoking	Yes 1 No 2 Not Applicable 8	
	Advice to start or do more physical activity	Yes 1 No 2	
248.	During the past 12 months have you visited to an AYUSH Practitioner for high blood pressure or hypertension?	Yes 1 No 2	If No, go to 250
249.	Are you currently taking any treatment/medicine from an AYUSH Practitioner for your high blood pressure?	Yes 1 No 2	
History of Diabetes			
250.	Has your blood sugar been measured in the last 12 months ?	Yes 1 No 2	
251.	Have you ever been told by a doctor or health worker that you have diabetes ?	Yes 1 No 2	If No, go to 253
252.	Are you currently receiving any of the following treatments/advice for diabetes prescribed by a doctor or other health worker as well as any advice ?		
	Insulin	Yes 1 No 2	
	Oral drug (medication that you have taken in the last 2 weeks).	Yes 1 No 2	
	Special Prescribed diet	Yes 1 No 2	
	Advice or treatment to lose weight	Yes 1 No 2	
	Advise to start or do more exercise	Yes 1 No 2	
253.	During the past 12 months have you visited/ seen an AYUSH Practitioner for diabetes?	Yes 1 No 2	If No, go to 301
254.	Are you currently taking any treatment/medicine from an AYUSH Practitioner for your diabetes?	Yes 1 No 2	

STEP 2. Physical Measurement			
Questions		Response	Skip
301.	Technician / Interviewer ID		
302.	Device ID for height and weight	Height..... <input type="text"/> <input type="text"/> Weight..... <input type="text"/> <input type="text"/>	
303.	Height	In Centimeter(cm)... <input type="text"/> <input type="text"/> <input type="text"/> . <input type="text"/>	
304.	Weight	In Kilograms (kg).... <input type="text"/> <input type="text"/> <input type="text"/> . <input type="text"/>	
305.	(For Women) Are you pregnant?	Yes 1 No 2	If Yes, go to 309
Waist Measurement			
306.	Device ID for waist	<input type="text"/> <input type="text"/>	
307.	Waist circumference Reading 1	In Centimeter (cm)... <input type="text"/> <input type="text"/> <input type="text"/> . <input type="text"/>	
308.	Waist circumference Reading 2	In Centimeter (cm)... <input type="text"/> <input type="text"/> <input type="text"/> . <input type="text"/>	
Blood Pressure and Pulse Rate			
309.	Technician ID	<input type="text"/> <input type="text"/>	
310.	Device ID for Blood Pressure	<input type="text"/> <input type="text"/>	
311.	Cuff Size Used	Small 1 Medium 2 Large 3	
312.	B.P. Reading 1	Systolic (mmHg) <input type="text"/> <input type="text"/> <input type="text"/> Diastolic (mmHg) <input type="text"/> <input type="text"/> <input type="text"/>	
313.	Pulse Rate Reading 1	<input type="text"/> <input type="text"/> <input type="text"/>	
314.	B.P. Reading 2	Systolic (mmHg) <input type="text"/> <input type="text"/> <input type="text"/> Diastolic (mmHg) <input type="text"/> <input type="text"/> <input type="text"/>	
315.	Pulse Rate Reading 2	<input type="text"/> <input type="text"/> <input type="text"/>	
316.	B. P. Reading 3	Systolic (mmHg) <input type="text"/> <input type="text"/> <input type="text"/> Diastolic (mmHg) <input type="text"/> <input type="text"/> <input type="text"/>	
317.	Pulse Rate Reading 3	<input type="text"/> <input type="text"/> <input type="text"/>	

Appendix - C

NATIONAL TECHNICAL ADVISORY COMMITTEE

Prof. N. K. Ganguly Former Director General, ICMR, New Delhi	Chair-person
Prof. K. S. Reddy President, Public Health Foundation of India, New Delhi	Member
Dr. R.L. Ichhpujani National Project Officer (IDSP), NICD, DGHS, MOHFW, New Delhi	Member
Dr. Rajesh Kumar Professor and Head, School of Public Health, PGI, Chandigarh	Member
Dr. K.R. Sunderam Professor & Head, Depart. of Biostatistics, Amrita Institute of Medical Sciences & Research Centre, Kochi, Kerala	Member
Prof. Arvind Pandey Director, National Institute of Medical Statistics, ICMR, New Delhi	Member
Dr. K. Anand Associate Professor, Centre for Community Medicine, AIIMS, New Delhi	Member
Dr. V. Mohan Director, Madras Diabetes Research Foundation, Chennai	Member
Dr. Bela Shah Scientist 'G' & Head - Division of Non-Communicable Diseases, ICMR, New Delhi	Member-Secretary

NATIONAL MONITORING COMMITTEE

Dr. Shiv Lal Spl. Director General, DGHS & Project Director (IDSP), NICD, MOHFW, New Delhi	Chair-person
Dr. S.K. Bhattacharya Addl. Director General, ICMR, New Delhi	Member
National Project Officer - IDSP, NICD, DGHS, MOHFW, New Delhi	Member
Dr. Bela Shah Scientist 'G' & Head - Division of Non-Communicable Diseases, ICMR, New Delhi	Member
Smt. Nirmala Kutty Under Secretary-Finance	Member
Dr. Shah Hossain CSU-IDSP, NICD, New Delhi	Member-Secretary

ICMR-PROJECT REVIEW GROUP

<p>Dr. L.M. Nath Consultant, New Delhi</p>	Chair-person
<p>Dr. Bela Shah Scientist 'G' & Head-Division of NCD, ICMR, New Delhi</p>	Member
<p>Prof. Arvind Pandey Director, NIMS, ICMR, New Delhi</p>	Member
<p>Dr. K. Anand Associate Professor, Centre for Community Medicine, AIIMS, New Delhi</p>	Member
<p>Dr. D. K. Shukla Scientist 'F', ICMR, New Delhi</p>	Member
<p>Dr. Prashant Mathur Scientist 'D', ICMR, New Delhi</p>	Member
<p>Dr. H. K. Chaturvedi Scientist 'E', NIMS, ICMR, New Delhi</p>	Member

WORLD BANK
(IDSP -NCD RISK FACTOR SURVEY)

Dr. G. Ramana
Mr. G.J. Sastry, Consultant

NATIONAL INSTITUTE OF COMMUNICABLE DISEASES, NEW DELHI
(CENTRAL SURVEILLANCE UNIT)

Dr. Shiv Lal
Spl. Director General, DGHS &
Project Director (IDSP)

Dr. D. Bachani
Dr. R.L. Ichhpujani
Dr. A. C. Dhariwal,
Dr. Shah Hossain
Dr. Pradeep Khasnobis

DIVISION OF NON-COMMUNICABLE DISEASES
INDIAN COUNCIL OF MEDICAL RESEARCH, NEW DELHI
PROJECT IMPLEMENTING AGENCY

Dr. Bela Shah, Scientist 'G' & Head

Dr. D. K. Shukla, Scientist 'F'
Dr. Prashant Mathur, Scientist 'D'

NATIONAL INSTITUTE OF MEDICAL STATISTICS, ICMR, NEW DELHI
NATIONAL NODAL AGENCY (NNA)

Prof. Arvind Pandey, Director
Dr. H. K. Chaturvedi, Scientist 'E'
Dr. Damodar Sahu, Scientist 'D'
Dr. Tulsi Adhikari, Scientist 'C'
Dr. Atul Juneja, Scientist 'C'
Mr. Kh. Jiten Kumar Singh, Scientist 'B'

Dr. B. N. Bhattacharya
Dr. Vinod K. Patro
Dr. Aikant Bhatti
Mr. Chandan Bandopadhyay
Dr. Vinod K. Upadhyay
Mr. Ashish Yadav
Mr. Sarad Mathur
Mr. Jeetendra Yadav
Mrs. Madhu Mehra
Mr. Charan Singh

NATIONAL INSTITUTE OF EPIDEMIOLOGY, CHENNAI

Regional Resource Centre

Prof. M.D. Gupte, Director

Dr. R. Rama Krishnan , Scientist 'E'

Dr. Prabhadeep Kaur, Scientist 'C'

Indian Institute of Health & Family Welfare, Hyderabad

State Survey Agency (SSA)

Dr. P. Satya Sekhar,
Principal Investigator

Dr. Prakash Bhatia, Co-Principal Investigator

Mr. M. Dharma Rao, Coordinator

Mr. Ch.V.S. Sita Rama Rao, Coordinator

Supervisor

Mr. M.Kiran Kumar

Ms. C.H.Padmarenuka

Mr. I.Ravi Kumar

Mr. S.K.Basha

Mr. M.Suresh Babu

Investigator

Ms. C. H. Renuka

Ms. G. Sobharani

Mr. P. Srinivasa

Mr. M. Hussaian

Mr. V. Narashimharaju

Mr. M. Ramesh Babu (Mapping, Listing)

Mr. C. H. Markandeya Prasad

Mr. K. Rajendraprasad

Mr. M. Ashok

Mr. B. Suresh

Mr. A. Satyadev

Mr. V. Mayachari

Mr. V. Lakshmi Pathi (Mapping, Listing)

Ms. V. L. Sailaja (F)

Mr. P. Bhaskar Reddy

Mr. B. Venkat Reddy

Mr. K. Srinivasyadv

Mr. V. Ravindar

Mr. U. Tarun (Mapping, Listing)

Ms. K.Adhilakshmi

Mr. P.Indrasena

Mr. S. Ramunaidu

Mr. K.Vasudevarao

Mr. N. Apparao

Mr. M. Rajendra (Mapping, Listing)

